

**CBE ID**

0053

**Title**

Osteoporosis Management in Women Who Had a Fracture

**Project**

Primary Care and Chronic Illness

**Endorsement Status**

Endorsement Removed

**E&M Committee Rationale/Justification**

Steward is no longer seeking endorsement.

**Is Under Review**

No

**Previous Endorsement Cycle**

Spring 2018

**Removal Date**

Wed, 04/16/2025 - 11:50

**Initial Endorsement**

Sun, 08/09/2009 - 20:00

**Steward**

National Committee for Quality Assurance

**1.0 New or Maintenance**

Maintenance

**1.1 Measure Structure**

Single Measure

**1.3 Electronic Clinical Quality Measure (eCQM)**

No

**1.6 Measure Description**

The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis.

**1.7 Measure Type**

Process

## 1.8 Level of Analysis

Clinician: Group/Practice, Clinician: Individual, Health Plan, Integrated Delivery System

## 1.9 Care Setting

Outpatient Services

## 1.14 Numerator

Patients who received either a bone mineral density test or a prescription for a drug to treat osteoporosis after a fracture occurs.

## 1.15 Denominator

Women who experienced a fracture, except fractures of the finger, toe, face or skull. Three denominator age strata are reported for this measure: Women age 50-64 Women age 65-85 Women age 50-85

## 1.20 Types of Data Sources

Claims Data, Electronic Health Data, Electronic Health Records: Electronic Health Records, Paper Patient Medical Records

## 6.1.2 Current or Planned Use(s)

Public Reporting, Quality Improvement with Benchmarking (external benchmarking to multiple organizations), Regulatory and Accreditation Programs

## 6.1.3 Current Use(s)

Public Reporting, Quality Improvement with Benchmarking (external benchmarking to multiple organizations), Regulatory and Accreditation Programs

## Exclusions

- Exclude women who had a bone mineral density test during the 24 months prior to the index fracture.
- Exclude women who had a claim/encounter for osteoporosis treatment during 12 months prior to the index fracture.
- Exclude women who received a dispensed prescription or had an active prescription to treat osteoporosis during the 12 months prior to the index fracture.
- Exclude women who are enrolled in a Medicare Institutional Special Needs Plan (I-SNP) or living long-term in an institution any time during the measurement year.
- Exclude women receiving hospice care during the measurement year.

## Measure Disclaimer

These performance Measures are not clinical guidelines and do not establish a standard of medical care, and have not been tested for all potential applications.

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY

KIND.

### **Risk Adjustment**

No risk adjustment or risk stratification

### **Target Population**

Elderly

### **Use In Federal Program**

Medicare Physician Quality Reporting System (PQRS), Physician Feedback/Quality and Resource Use Reports (QRUR), Physician Value-Based Payment Modifier (VBM)

### **Steward Organization**

National Committee for Quality Assurance

### **Steward POC email**

[measureendorsement@ncqa.org](mailto:measureendorsement@ncqa.org)