

2024 Measure Set Review (MSR): Final Preliminary Assessment

The following information was sourced in June of 2024 from the Centers for Medicare & Medicaid Services Measures Inventory Tool (CMIT), discussions with CMS program leads, and publicly available CMS datasets (see links below).

Measure Information

CMIT ID	Title
00575-01-C-IRFQR	Potentially Preventable 30-Day Post-Discharge Readmission Measure for Inpatient Rehabilitation Facility Quality Reporting Program
Measure Steward	CMS Program
Centers for Medicare & Medicaid Services (CMS)	Inpatient Rehabilitation Facility Quality Reporting

Measure Overview

Rationale: This measure is part of potentially preventable readmission (PPR) measures for post-acute care (PAC) that estimate the risk-standardized rate of unplanned, potentially preventable readmissions for patients (Medicare fee-for-service [FFS] beneficiaries) who receive services in one of the following post-acute care provider types: skilled nursing facilities (SNFs), inpatient rehabilitation facilities (IRFs), and long-term care hospitals (LTCH).

Description: This measure is conceptualized uniformly across the PAC settings, in terms of the definition of the PPR outcome, the approach to risk adjustment, and the measure calculation. These outcome measures reflect readmission rates for patients who are readmitted to a short-stay acute-care hospital or an LTCH with a principal diagnosis considered to be unplanned and potentially preventable.

Numerator: The numerator is mathematically related to the number of patients in the target population who have the event of a potentially preventable readmission in the 30-day post-discharge window. The measure does not have a simple form for the numerator and denominator; that is, the risk-adjustment method used does not make the observed number of potentially preventable readmissions the numerator and a predicted number the denominator. Instead, the numerator is the risk-adjusted estimate of the number of potentially preventable readmissions that occurred within 30 days from discharge. This estimate includes risk adjustment for patient characteristics and a statistical estimate of the facility effect beyond patient mix.

Denominator: The risk-adjusted expected number of potentially preventable readmissions. This estimate includes risk adjustment for patient characteristics with the facility effect removed. The expected number of potentially preventable readmissions is the predicted number of risk-adjusted potentially preventable readmissions if the same patients were treated at the average PAC provider appropriate to the measure.

Exclusions: 1) Patients who died during the IRF stay. 2) Patients/residents less than 18 years old. 3) Patients who were transferred at the end of a stay to another IRF, LTCH, or short-term acute care hospital. 4) Patients/residents who were not continuously enrolled in Part A FFS Medicare for the 12 months prior to the IRF admission date and at least 30 days after IRF discharge date. 5) Patients who did not have a short-term acute-care stay within 30 days prior to an IRF admission date. 6) Patients discharged against medical advice (AMA).



7) Patients/residents for whom the prior short-term acute-care stay was for nonsurgical treatment of cancer. 8) Patients who were transferred to a federal hospital from the PAC facility. 9) Patients who received care from a provider located outside of the United States, Puerto Rico, or a U.S. territory.

or a U.S. territory.	
Measure type: Outcome	Measure is a composite: No Measure is digital and/or an eCQM: No
Level(s) of analysis/measured entity: Facility/Hospital/Agency	Care setting:
Risk adjustment and/or stratification: Yes, risk adjustment includes variables such as age, sex, category of principal diagnosis from the prior proximal acute stay and comorbidities. More information on IRFQR program risk adjustment can be found here .	Data source(s): Claims Data
Data collection method: Claims data review	Reporting frequency: Annually
All required data are collected as part of clinical workflow: Yes, routinely collected in claims data.	Reporting overlap with similar/related measures: Yes, part of four measure variations assessing 30-day post-discharge readmission across PAC/LTC settings. Includes 00575-02-C-LTCHQR, 00575-01-C-IRFQR, 00575-03-C-SNFQRP, and 00575-04-
	C-HHQR.

Measure Status	
Current CBE Endorsement Status: Not Endorsed	CBE Endorsement History: None

II. Measure Performance

00575-01-C-IRFQR Performance in 2020-2022

For this measure, the MSR evaluation and analysis team reviewed the publicly available dataset Inpatient Rehabilitation Facility - Provider Data and archived Inpatient Rehabilitation Facilities.

Figure 1 is a boxplot that shows the distribution of the performance over the past 3 years (where available). For each performance year, the dots indicate the lower 5th and upper 95th percentiles, and the vertical line is the range between these values (90% of the measure scores



are between the dots). The box spans the lower 25th to the upper 75th percentile (50% of the measure scores are within the box). The horizontal line in the box indicates the median score, and the "+" indicates the mean score. This plot can be used to assess overall trends in the score over time.

Interpretation of measure performance scores: In the plot below, the median score increases substantially from about 6.8 in 2020-2021 to about 8.5 in 2022. This suggests that there was improvement across entities from 2021 to 2022.

Figure 1. Boxplot of Measure Score by Year

Importance Table

This table shows the relative spread of the scores and how many patients are impacted. Often the lowest or highest deciles (which, by definition, each represent 10% of the entities) may represent a disproportionately higher or lower percentage of patients. If the lowest decile contains only 5% of the patients for example, it suggests that low patient population may be related to low scores.

Interpretation of measure performance scores: The table can also be used to evaluate the impact of improving the score. It is common practice to use the performance of the top 20% of the entities as a benchmark. Here, 20% of the entities perform better than the 3rd Decile (8.2), which could be considered the benchmark. The number of adverse events for each decile can be estimated by multiplying the total patients by the corresponding rate. Here the estimated total number of adverse events across all deciles is 47,716. If Deciles 4-10 performed at the benchmark of 8.2, there would be an estimated 8% fewer adverse events (about 43,762).



Table 1. Importance (Decile by measure score, 2022)

Data Type	Overall	Min	Decile 1	Decile 2	Decile 3	Decile 4	Decile 5	Decile 6	Decile 7	Decile 8	Decile 9	Decile 10	Max
Mean Score (SD)	8.63 (0.80)	6.2	7.4	7.9	8.2	8.3	8.5	8.7	8.9	9.1	9.5	10.3	12.3
Entities	1,069	1	107	107	107	107	107	107	107	107	107	106	1
Total Patients	541,632	2,269	63,869	47,059	47,472	42,622	39,581	41,732	47,710	53,347	69,087	89,153	1,085

Reliability Tables

Two tables are used to summarize reliability. For Table 2, entities are sorted by patient volume, and the average reliability is reported along with the number of entities and average number and total patients for each decile. These tables can be used to assess the impact of population size on the reliability of an entity's measure score. In cases where reliability has a strong relationship to population size, reliability will be the lowest at Decile 1 and progressively increase up to Decile 10.

For Table 3, entities are sorted by reliability, and the average reliability by decile is reported. Mean, standard deviation, minimum and maximum reliability, and inter-quartile range (IQR) are also included. This table can be used to see the distribution of the reliability of the entities. A measure score is generally considered reliable when the reliability for at least 70% of the individual entities is above 60%.

Table 2. Reliability (Decile by denominator – target population size)

(based on observed values)

Data Type	Overall	Min	Decile 1	Decile 2	Decile 3	Decile 4	Decile 5	Decile 6	Decile 7	Decile 8	Decile 9	Decile 10	Max
Mean Target Population Size	507	27	74	139	190	236	293	377	491	674	966	1,636	4,567
Mean Reliability	44.0	30.5	32.4	35.4	36.4	38.4	39.6	42.8	45.1	50.1	55.8	64.0	80.9
Entities	1,069	1	107	107	107	107	107	107	107	107	107	106	1
Total Patients	541,632	27	7,963	14,912	20,313	25,232	31,350	40,362	52,546	72,155	103,349	173,450	4,567



Table 3. Mean reliability (By reliability decile)

Mean	SD	Min	Decile 1	Decile 2	Decile 3	Decile 4	Decile 5	Decile 6	Decile 7	Decile 8	Decile 9	Decile 10	Max	IQR
44.0	10.2	25.0	30.5	34.1	36.3	38.4	40.5	42.8	46.0	50.7	55.8	65.0	80.9	14.3

Interpretation: The overall variation between entities (as estimated by the variance of the measure scores) is somewhat low relative to the variation within each entity. Over 90% of the entities have an estimated reliability of less than 60%, suggesting that this measure may not be effective in differentiating entities by quality of performance.