

## THE FIM® INSTRUMENT

The FIM® Instrument was intended to measure a patient's disability in terms of the need for assistance. Facilities world-wide use the FIM® Instrument to document patient functionality throughout medical rehabilitation, and the Centers for Medicare and Medicaid Services (CMS) have adopted it for the Inpatient Rehabilitation Facility – Prospective Payment System (IRF-PPS). Since its introduction, over 960 articles of supporting research and study have been published.

## REFERENCES

1. Abrahamson SJ, Khan F. Brief osteoporosis education in an inpatient rehabilitation setting improved knowledge of osteoporosis in elderly patients with low-trauma fractures. *Int J Rehabil Res.* 2006;29(1):61-64.
2. Acevedo-Navarro L, Alvelo J, Torres B. Activities of daily functioning and community adjustment of veterans in the VA residential care program. *P R Health Sci J.* 1992;11(1):33-41.
3. Adamovich BLB. Pitfalls in functional assessment: a comparison of FIM ratings by speech language pathologists and nurses. *Neurorehabilitation.* 1992;2(4):42-51.
4. Adunsky A, Arad M, Levi R, Blankstein A, Zelig G, Mizrahi E. Five-year experience with the 'Sheba' model of comprehensive orthogeriatric care for elderly hip fracture patients. *Disabil Rehabil.* 2005;27(18-19):1123-1127.
5. Adunsky A, Fleissig Y, Levenkrohn S, Arad M, Noy S. Clock drawing task, mini-mental state examination and cognitive-functional independence measure: relation to functional outcome of stroke patients. *Arch Gerontol Geriatr.* 2002;35(2):153-160.
6. Adunsky A, Fleissig Y, Levenkrohn S, Arad M, Noy S. A comparative study of mini-mental test, clock drawing task, and cognitive-FIM in evaluation function outcome of elderly hip patients. *Clin Rehabil.* 2002;16(4):414-419.
7. Adunsky A, Levenkrohn S, Fleissig Y, Arad M, Heruti RJ. Rehabilitation outcomes in patients with full weight-bearing hip fractures. *Arch Gerontol Geriatr.* 2001;33(2):123-131.
8. Adunsky A, Levenkrohn S, Fleissig Y, Chetrit A, Blumstein Z. [Evaluation of predictive factors for stroke rehabilitation.] *Harefuah.* 1998;153(12):569-573,656.
9. Adunsky A, Levenkrohn S, Fleissig Y, Blumstein Z, Chetrit A. In-hospital referral source and rehabilitation outcome of elderly stroke patients. *Aging (Milano).* 2001;13(6):430-436.

10. Adunsky A, Levy R, Heim M, Mizrahi E, Arad M. The unfavorable nature of preoperative delirium in elderly hip fractured patients. *Arch Gerontol Geriatr.* 2003;36(1):67-74.
11. Adunsky A, Lasky A, Arad M, Heruti RJ. A comparative study of rehabilitation outcomes of elderly hip fracture patients: the advantage of a comprehensive orthogeriatric approach. *J Gerontol A Biol Sci Med Sci.* 2003;58(6):542-547.
12. Agarwal V, Mcrae MP, Bhardwaj A, Teasell RW. A model to aid in the prediction of discharge location for stroke rehabilitation patients. *Arch Phys Med Rehabil.* 2003;84(11):1703-1709.
13. Aisen ML, Sevilla D, Gibson G, Kutt H, Blau A, Edelstein L, Hatch J, Blass J. A double-blind, placebo-controlled study of 3,4-diaminopyridine as a treatment for amyotrophic lateral sclerosis. *Journal of the Neurological Sciences.* 1995;129(1):21-24.
14. Aitken ME, Jaffe KM, DiScala C, Rivara FP. Functional outcome in children with multiple trauma without significant head injury. *Arch Phys Med Rehabil.* 1999;80(8):889-895.
15. Aitken ME, Tilford JM, Barrett KW, et al. Health status of children after admission for injury. *Pediatrics.* 2002;110(2):337-342.
16. Aito S, D'Andrea M, Werhagen L, et al. Neurological and functional outcome in traumatic central cord syndrome. *Spinal Cord.* 2007;45(4):292-7.
17. Akmal M, Trivedi R, Sutcliffe J. Functional outcome in trauma patients with spinal injury. *Spine.* 2003;28(2):180-185.
18. Alexander DN, Cen S, Sullivan KJ, Bhavnani G, Ma X, Azen SP. Effects of acupuncture treatment on poststroke motor recovery and physical function: a pilot study. *Neurorehabil Neural Repair.* 2004;18(4):259-267.
19. Alexander MP. Stroke rehabilitation outcome: a potential use of predictive variables to establish levels of care. *Stroke.* 1994;25(1):128-134.
20. Alexander MP. Chronic akinetic mutism after mesencephalic-diencephalic infarction: remediated with dopaminergic medications. *Neurorehabil Neural Repair.* 2001;15(2):151-156.
21. Alfred WG, Fuhrer MJ, Rossi CD. Vocational development following severe spinal cord injury: a longitudinal study. *Archives of Physical Medicine & Rehabilitation.* 1987; 68(12):854-857.
22. Allen BP, Agha Z, Duthie EH, Layde PM. Minor depression and rehabilitation outcome for older adults in subacute care. *J Behav Health Serv Res.* 2004;31(2):189-198.
23. Alt-White AC, Romano EL. An interdisciplinary approach to improving the quality of life for nursing home residents. *Nursing Connections.* 1993;6(4):51-59.

24. Amman P, Herter-Clavel C, Lubrano A, Rizzoli R. A single bisphosphonate infusion is associated with improved functional capacity in elderly subjects with primary hyperparathyroidism. *Aging Clin Exp Res*. 2003;15(6):500-504.
25. Amory JK, Chansky HA, Chansky KL, et al. Preoperative supraphysiological testosterone in older men undergoing knee replacement surgery. *J Am Geriatr Soc*. 2002;50(10):1698-1701.
26. Amsters DI, Pershouse KJ, Price GL, Kendall MB. Long duration spinal cord injury: perceptions of functional change over time. *Disabil Rehabil*. 2005;27(9):489-497.
27. Ancheta J, Husband M, Law D, Reding M. Initial functional independence measure score and interval post stroke help assess outcome, length of hospitalization, and quality of care. *Neurorehabil Neural Repair*. 2000;14(2):127-134.
28. Anderson CJ, Krajci KA, Vogel LC. Life satisfaction in adults with pediatric-onset spinal cord injuries. *J Spinal Cord Med*. 2002;25(3):184-190.
29. Anderson CJ, Vogel LC. Employment outcomes of adults who sustained spinal cord injuries as children or adolescents. *Arch Phys Med Rehabil*. 2002;83(6):791-801.
30. Anderson CW. The Western Physical Performance Analysis: a functional outcome tool for the severely impaired neurological patient. *Neurology Report*. 1993;17(4):35-38.
31. Anderson K, Aito S, Atkins M, et al.; Functional Recovery Outcome Measures Work Group. Functional recovery measures for spinal cord injury: an evidence-based review for clinical practice and research. *J Spinal Cord Med*. 2008;31(2):133-44.
32. Andren E, Grimby G. Dependence and perceived difficulty in activities of daily living in adults with cerebral palsy and spinal bifida. *Disabil Rehabil*. 2000;22(7):299-307.
33. Andren E, Grimby G. Andre E, Grimby G. Dependence in daily activities and life satisfaction in adult subjects with cerebral palsy or spinal bifida; a follow-up study. *Disabil Rehabil*. 2004;26(9):528-536.
34. Andren E, Grimby G. Activity limitations in personal, domestic and vocational tasks: a study of adults with inborn and early acquired mobility disorders. *Disabil Rehabil*. 2004;26(50):262-271.
35. Andrews AW, Bohannon RW. Discharge function and length of stay for patients with stroke are predicted by lower extremity muscle force on admission to rehabilitation, *Neurorehab Neural Repair*. 2001;15(2):93-97.
36. Ang YH, Chan DK, Heng DM, Shen Q. Patient outcomes and length of stay in a stroke unit offering both acute and rehabilitation services. *Med J Aust*. 2003;178(7):333-336.
37. Appelros P, Nydevik I. [For whom is training after stroke most beneficial? Selection method exists and should be used]. *Lakartidningen*. 1999;96(19):2343-2347.
38. Appelros P, Nydevik I, Karlsson GM, Thornwalls A, Seiger A. Recovery from unilateral neglect after right-hemisphere stroke. *Disabil Rehabil*. 2004;26(8):471-477.

39. Arad M, Abboud S, Radi MM, Adunsky A. Heart rate variability parameters correlate with functional independence measures in ischemic stroke patients. *J Electrocardiol.* 2002;35(Suppl):243-246.
40. Arad M, Ring H, Abboud S, Radai MM, Tamir A, Adunsky A. Dynamic cardiophysiologic variables correlate with lesion location and FIM in patients with ischemic stroke. *Am J Phys Med Rehabil.* 2002;81(8):590-596.
41. Aras MD, Kaya A, Cakc A, Gokkaya KO. Functional outcome following traumatic brain injury: the Turkish experience. *Int J Rehabil Res.* 2004;27(4):257-260.
42. Azaula M, Msall ME, Buck G, Tremont MR, Wilczenski F, Rogers BT. Measuring functional status and family support in older school-ages children with cerebral palsy: comparison of three instruments. *Arch Phys Med Rehabil.* 2000;81(3):307-311.
43. Azouvi P, Mane M, Thiebaut JB, Denys P, Remyneris O, Bussel B. Intrathecal baclofen administration for control of severe spinal spasticity: functional improvement and long-term follow-up. *Archives of Physical Medicine & Rehabilitation.* 1996;77(1):35-9.
44. Bagg S, Pombo AP, Hopman W. Effect of age on functional outcomes after stroke rehabilitation. *Stroke.* 2002;33(1):179-185.
45. Baguley IJ, Nicholls JL, Felmingham KL, Crooks J, Gurka JA, Wade LD. Dysautonomia after traumatic brain injury: a forgotten syndrome? *J Neurol Neurosurg Psychiatry.* 1999;67(1):39-43.
46. Baker JG, Granger CV, Fielder RC. A brief outpatient functional assessment measure: validity using Rasch measures. *American Journal of Physical Medicine and Rehabilitation.* 1997;76(1):8-13.
47. Baker JG, Granger CV, Ottenbacher KJ. Validity of a brief outpatient functional assessment measure. *American Journal of Physical Medicine and Rehabilitation.* 1996;75(5):356-363.
48. Bakheit AMO, Harries SR, Hull RG. Validity of a self-administered version of the Barthel Index in patients with rheumatoid arthritis. *Clinical Rehabilitation.* 1995;9(3):234-7.
49. Bakhshi V, Elliot M, Gentili A, Godschalk M, Mulligan T. Testosterone improves rehabilitation outcomes in older men. *J Am Geriatric Soc.* 2004;48(5):550-553.
50. Baldrige RB. Functional assessment of measurements. *Neurology Report.* 1993;17(4):3-10.
51. Baldry Currrens JA. Evaluation of disability and handicap following injury. *Injury.* 2000;31(3):99-106.
52. Baldry Currrens JA, Coats TJ. The timing of disability measurements following injury. *Injury.* 2000;31(2):93-98.

53. Ballinger DA, Rintala DH, Hart KA. The relation of shoulder pain and range-of-motion problems to functional limitations, disability, and perceived health of men with spinal cord injury: a multifaceted longitudinal study. *Arch Phys Med Rehabil.* 2000;81(12):1575-1581.
54. Barnes C, Conner D, Legault L, Reznickova N, Harrison-Felix C. Rehabilitation outcomes in cognitively impaired patients admitted to skilled nursing facilities from the community. *Arch Phys Med Rehabil.* 2004;85(10):1602-1607.
55. Barnes MP, Kent RM, Semlyen JK, McMullen KM. Spasticity in multiple sclerosis. *Neurorehab Neural Repair.* 2003;17(1):66-70.
56. Bates BE, Stineman MG. Outcome indicators for stroke: application of an algorithm treatment across the continuum of postacute rehabilitation services. *Arch Phys Med Rehabil.* 2000;81(11):1468-1478.
57. Beck LA, Harris MR, Basford J. Factors influencing functional outcome and discharge disposition after thoracic spinal cord injury. *SCI Nurs.* 1999;16(4):172-132.
58. BeDell KK, Scremin AME, Perell KL, Kunkel CF. Effects of functional electrical stimulation-induced lower extremity cycling on bone density of spinal cord-injured patients. *American Journal of Physical Medicine and Rehabilitation.* 1996;75:29-34.
59. Bell KR, Tallman CA. Community re-entry of long-term institutionalized brain-injured persons. *Brain Injury.* 1995;9(3):315-320.
60. Bell KR, Temkin NR, Esselman PC, et al. The effect of a scheduled telephone intervention on outcome after moderate to severe traumatic brain injury: a randomized trial. *Arch Phys Med Rehabil.* 2005;86(5):851-856.
61. Beloosesky Y, Grinblat J, Epelboym B, Hendel D. Dementia does not significantly affect complications and functional gain in elderly patients operated on for intracapsular hip fracture. *Arch Orthop Trauma Surg.* 2001;121(5):257-260.
62. Benaim C, Perennou DA, Villy J, Rosseaux M, Pelissier JY. Validation of a standardized assessment of postural control in stroke patients: the Postural Assessment Scale for Stroke Patients (PASS). *Stroke.* 1999;30(9):1862-1868.
63. Beninato M, O'Kane KS, Sullivan PE. Relationship between motor FIM and muscle strength in lower cervical-level spinal cord injuries. *Spinal Cord.* 2004;42(9): 533-540.
64. Beninato M, Gil-Body LM, Salles S, Stark PC, Black-Schaffer RM, Stein J. Determination of the minimal clinically important difference in the FIM instrument in patients with stroke. *Arch Phys Med Rehabil.* 2006;87(1):32-29.
65. Berner YN, Lif Kimchi O, Karpin H, Finkelrov B. Triple task Clock Completion Test (CCT) as a predictor of functional outcome in geriatric rehabilitation. *Arch Gerontol Geriatr.* 2004;39(2):117-124.
66. Berner YN, Lif Kimchi O, Spokoyny V, Finkelrov B. The effect of electric stimulation treatment on the functional rehabilitation of acute geriatric patients with stroke: a preliminary study. *Arch Gerontol Geriatr.* 2004;39(2):125-132.

67. Berry JF, Hitzman S, Stewart GW, Darwin P. A survey of attendant care arrangements in indigent persons with spinal cord injury. *SCI Psychosocial Process*. 1995;8(3):112-117.
68. Bethoux F, Calmels P, Gautheron V, Minaire P. Quality of life of the spouses of stroke patients: a preliminary study. *Int J Rehabil Res*. 1996;19(4):291-299.
69. Bezner JR, Hunter DL. Wellness perception in persons with traumatic brain injury and its relation to functional independence. *Arch Phys Med Rehabil*. 2001;82(6):787-792.
70. Bhandari VK, Kushel M, Price L, Schillenger D. Racial disparities in outcomes of inpatient stroke rehabilitation. *Arch Phys Med Rehabil*. 2005;86(11):2081-2086.
71. Black K, Zafonte R, Millis S, et al. Sitting balance following brain injury: does it predict outcome? *Brain Inj*. 2000;14(2):141-152.
72. Black TM, Soltis T, Bartlett C. Using the Functional Independence Measure instrument to predict stroke rehabilitation outcomes. *Rehabi Nurs*. 1999;24(3):109-114, 121.
73. Blackerby W, King J. The effects of the interdisciplinary team on the rehabilitation process and outcome. *Archives of Physical Medicine & Rehabilitation*. 1992;73(10):977.
74. Bluvstein V, Front L, Itzkovich M, et al. A new grading for easy and concise description of functional status after spinal cord lesions. *Spinal Cord*. 2011; Aug 2. doi: 10.1038/sc.2011.84. [Epub ahead of print]
75. Bode RK, Heinemann AW. Course of functional improvement after stroke, spinal cord injury, and traumatic brain injury. *Arch Phys Med Rehabil*. 2002;83(1):100-106.
76. Bogner JA, Corrigan JD, Fugate L, Mysiw WJ, Clinchot D. Role of agitation in prediction of outcomes after traumatic brain injury. *Am J Phys Med Rehabil*. 2001;80(9):636-644.
77. Bohannon RW, Ahlquist M, Lee N, Maljanian R. Functional gains during acute hospitalization for stroke. *Neurorehabil Neural Repair*. 2003;17(3):192-195.
78. Bohannon RW, Leary KM. Standing balance and function over the course of acute rehabilitation. *Archive of Physical Medicine and Rehabilitation*. 1995;76(11):1994-1996.
79. Bohannon RW. Standing balance, lower extremity muscle strength, and walking performance of patients referred for physical therapy. *Perceptual & Motor Skills*. 1995; 80(2):379-385.
80. Bohannon, RW. Walking after stroke: comfortable versus maximum safe speed. *International Journal of Rehabilitation Research*. 1992;15:246-248.
81. Bohannon RW, Cooper J. Total knee arthroplasty: evaluation of an acute care rehabilitation program. *Archives of Physical Medicine & Rehabilitation*. 1993;74(10):1091-1094.

82. Bohannon RW, Kloter KS, Cooper JA. Outcome of patient with hip fracture treated by physical therapy in an acute hospital. *Topics in Geriatric Rehabilitation*. 1990;6(2):51-58.
83. Bohannon RW, Leary KM, Cooper J. Independence in floor-to-stand transfers soon after stroke. *Topics in Geriatric Rehabilitation*. 1995;11(1):6-9.
84. Bombardier CH, Stroud MW, Esselman PC, Rimmele CT. Do preinjury alcohol problems predict poorer rehabilitation progress in persons with spinal cord injury? *Arch Phys Med Rehabil*. 2004;85(9):1488-1492.
85. Bonan IV, Guettard E, Leman MC, Colle FM, Yelnik AP. Subjective visual vertical perception relates to balance in acute stroke. *Arch Phys Med Rehabil*. 2006;87(5):642-646.
86. Bonaroti D, Akers JM, Smith TM, Milcahey MJ, Betz RR. Comparison of functional electrical stimulation to long leg braces for upright mobility for children with complete thoracic level spinal injuries. *Arch Phys Med Rehabil*. 1999;80(9):1047-1053.
87. Bondanelli M, Ambrosio MR, Margutti A, et al. Evidence for integrity of the growth hormone/insulin-like growth factor-1 axis in patients with severe head trauma during rehabilitation. *Metabolism*. 2002;51(10):1363-1369.
88. Bondanelli M, Ambrosio MR, Onofri A, et al. Predictive value of circulating IGF-I levels in ischemic stroke outcome. *J Clin Endocrinol Metab*. 2006;91(10):3928-34.
89. Borgel F. [Rehabilitation management in neurologic follow-up of the multiple sclerosis patient]. *Rev Neurol (Paris)*. 2001;157(8-9):1135-1138.
90. Bottemiller KL, Bieber PL, Basford JR, Harris M. FIM score, FIM efficiency, and discharge disposition following inpatient stroke rehabilitation. *Rehabil Nurs*. 2006;31(1):22-25.
91. Bottger S, Prosiegel M, Steiger HJ, Yassouridis A. Neurobehavioural disturbances, rehabilitation outcome, and lesion site in patients after rupture and repair of anterior communicating artery aneurysm. *J Neurol Neurosurg Psychiatry*. 1998;65(1):93-102.
92. Boyd JA. Assessing functional ability and the team approach in the home. *Journal of Home Care Practice*. 1988;1(1):11-24.
93. Brachner A. [Function independence measure (FIM)]. *Osterr Krankenpflegez*. 1998;51(8-9):20.
94. Bracken MB, Holford TR. Neurological and functional status 1 year after acute spinal cord injury: estimates of functional recovery in National Acute Spinal Cord Injury Study II from results modeled in National Acute Spinal Cord Injury Study III. *J Neurosurg*. 2002;96(3 Suppl):259-266.
95. Bracken MB, Shepard MJ, Holford TR, et al. Administration of methylprednisolone for 24 or 48 hours or tirilazad mesylate for 48 hours in the treatment of spinal cord injury. Results of the Third National Acute Spinal Cord Injury Randomized Controlled Trial. National Acute Spinal Cord Injury Study. *JAMA*. 1997;277(20):1597-1604.

96. Brautigam K, Flemming A, Schultz H, Dassen T. [How reliable is the Functional Independence Measure (FIM)?]. *Pflege*. 2002;15(3):131-136.
97. Brock KA, Goldie PA, Greenwood KM. Evaluating the effectiveness of stroke rehabilitation: choosing a discriminative measure. *Arch Phys Med Rehabil*. 2002;83(1):92-99.
98. Brooks D, Davis AM, Nagile G. Validity of 3 physical performance measures in inpatient geriatric rehabilitation. *Arch Phys Med Rehabil*. 2006;87(1):105-110.
99. Brosseau L, Philippe P, Boulanger YL. The Functional Independence Measure: validity of selected assessment methods applied to stroke survivors. *Topics in Geriatric Rehabilitation*. 1995;11(1):75-86.
100. Brosseau L, Potvin L, Philippe P, Boulanger YL, Dutil E. The construct validity of the Functional Independence Measure as applied to stroke patients. *Physiotherapy Theory and Practice*. 1996;12(3):161-171.
101. Brosseau L, Wolfson C. The inter-rater reliability and construct validity of the Functional Independence Measure for multiple sclerosis subjects. *Clinical Rehabilitation*. 1994;8(2):107-115.
102. Buchanan JL, Andres PL, Haley SM, Paddock SM, Zaslavsky AM. An assessment tool translation study. *Health Care Financ Rev*. 2003;24(3):45-60.
103. Buchanan JL, Andres PL, Haley SM, Paddock SM, Zaslavsky AM. Evaluating the planned substitution of the minimum dataset-post acute care for use in the rehabilitation hospital prospective payment system. *Med Care*. 2004;42(2):155-163.
104. Buia JM, Carlier P, Belkacemi A. Interest in the study of the Functional Independence Measure (FIM) in a geriatric rehabilitation unit. *Annales de Readaptation et de Medecine Physique*. 1995;38(7):435-41.
105. Bunch WH, Dvorchak VM. The "value" of Functional Independence Measure scores. *American Journal of Physical Medicine and Rehabilitation*. 1994;73(1): 40-43.
106. Burke DT, Shah MK, Dorvlo AS, Al-Adawi S. Rehabilitation outcomes of cardiac and non-cardiac anoxic brain injury: a single institution experience. *Brain Inj*. 2005;19(9):675-680.
107. Burnett DM, Kolakowsky-Hayner SA, Slater D, et al. Ethnographic analysis of traumatic brain injury in patients in the national Model Systems database. *Arch Phys Med Rehabil*. 2003;84(2):263-267.
108. Burnett DM, Silver TM, Kolakowsky-Hayner SA, Cifu DX. Functional outcome for African Americans and Hispanics treated at a traumatic brain injury model systems centre. *Brain Inj*. 2000;14(8):713-718.
109. Bushnik T, Hanks RA, Kreutzer J, Rosenthal M. Etiology of traumatic brain injury: characterization of differential outcomes up to 1 year postinjury. *Arch Phys Med Rehabil*. 2003;84(2):255-262.



110. Byrnes MB, Powers FF. FIM: its use in identifying rehabilitation needs in the head-injured patient. *Journal of Neuroscience Nursing*. 1989;21(1):61-63.
111. Caban P. A new approach to the assessment of aged community service users. *Australian Journal of Rural Health*. 1993;1(4):31-37.
112. Calmels P, Vedel E, Bethous F, Charmet E, Minaire P. [The Functional Independence Measure (FIM): interest of the telephone administration]. *Annales de Readaptation et de Médecine Physique*. 1994;37:469-476.
113. Calmels P, Pereira A, Domenach M, Pallot Prades B, Alexandre C, Minaire P. Functional ability and quality of life in rheumatoid arthritis: evaluation using the Functional Independence Measure and the reintegration to normal living index. *Revue du Rhumatisme English Edition*. 1994;61(11):723-31.
114. Calmels P, Ebermeyer E, Bethoux F, Gonard C, Fayolle-Minon I. [Relationship between burden of care at home and functional independence measure after stroke]. *Ann Readapt Med Phys*. 2002;45(3):105-113.
115. Calvanio R, Burke DT, Kim HJ, et al. Naltrexone: effects on motor function, speech, and activities of daily living in a patient with traumatic brain injury. *Brain Inj*. 2000;14(10):933-942.
116. Calvanio R, Williams R, Burke DT, et al. Acquired brain injury, visual attention, and the useful field of view test: a pilot study. *Arch Phys Med Rehabil*. 2004;85(3):474-478.
117. Cantalloube S, Monteil I, Lamotte D, Mailhan L, Thoumie P. [Strength, postural and gait changes following rehabilitation in multiple sclerosis: a preliminary study]. *Ann Readapt Med Phys*. 2006;49(4):143-149.
118. Caplan GA, Coconis J, Board N, Sayers A, Woods J. Does home treatment affect delirium? A randomized controlled trial of rehabilitation of elderly and care at home or usual treatment (The REACH-OUT trial). *Age Aging*. 2006;35(1):53-60.
119. Cardenas DD, Hoffman JM, Kirshblum S, McKinley W. Etiology and incidence of rehospitalization after traumatic spinal cord injury: a multicenter analysis. *Arch Phys Med Rehabil*. 2004;85(11):1757-1763.
120. Carey, R. Integrating case management, program evaluation, and marketing for inpatient and outpatient rehabilitation programs. *Advances in Clinical Rehabilitation*. 1990:219-249.
121. Carlson JE, Zocchi KA, Bettencourt DM, et al. Measuring frailty in the hospitalized elderly: concept of functional homeostasis. *Am J Phys Med Rehabil*. 1998;77(3):252-257.
122. Carod-Artel FJ, Medeiros MS, Horan TA, Braga LW. Predictive factors of functional gain in long-term stroke survivors admitted to a rehabilitation programme. *Brain Inj*. 2005;19(9):667-673.
123. Carter GT, Weiss MD, Lou JS, et al. Modafinil to treat fatigue in amyotrophic lateral sclerosis: an open label pilot study. *Am J Hosp Palliat Care*. 2005;22(1):55-59.

124. Carter GM, Relles DA, Ridgeway GK, Rimes CM. Measuring function for Medicare inpatient rehabilitation payment. *Health Care Financing Review*. 2003;24(3):25-43.
125. Casino CM, Blando AV. Calcification of ligamentum flavum causing spinal cord compression in a stroke patient. *Archives of Physical Medicine & Rehabilitation*. 1994;75(2):226-228.
126. Catz A, Itzkovich M, Agranov E, Ring H, Tamir A. SCIM—Spinal Cord Independence Measure: a new disability scale for patients with spinal cord lesions. *Spinal Cord*. 1997;35(12):850-856.
127. Catz A, Itzkovich M, Aragnov E, Ring H, Tamir A. The spinal cord independence measure (SCIM): sensitivity to functional changes in sub groups of spinal cord lesion patients. *Spinal Cord*. 2001;39(2):97-100.
128. Catz A, Itzkovich M, Steinberg F, et al. The Catz-Itzkovich SCIM: a revised version of the Spinal Cord Independence Measure. *Disabil Rehabil*. 2001;23(6):263-268.
129. Catz A, Itzkovich M, Steinberg F, et al. Disability assessment by a single rater or a team: a comparative study with the Catz-Itzkovich spinal cord independence measure. *J Rehabil Med*. 2002;34(5):226-230.
130. Catz A, Itzkovich M, Tamir A, et al. [SCIM—spinal cord independence measure (version II): sensitivity to functional changes]. *Harefuah*. 2002;141(12):1025-1031,1091.
131. Cavanagh SJ, Hogan K, Gordon V, Fairfax J. Stroke-specific FIM models in an urban population. *J Neurosci Nurs*. 2000;32(1):17-21.
132. Chae J, Johnston M, Kim H, Zorowitz R. Admission motor impairment as a predictor of physical disability after stroke rehabilitation. *American Journal of Physical Medicine and Rehabilitation*. 1995;(3):218-223.
133. Chae J, Zorowitz R. Functional status of cortical and subcortical nonhemorrhagic stroke survivors and the effect of lesion laterality. *Am J Phys Med Rehabil*. 1998;77(5):415-420.
134. Chae J, Zorowitz RD, Johnston MV. Functional outcome of hemorrhagic and nonhemorrhagic stroke patients after in-patient rehabilitation. *Am J Phys Med Rehabil*. 1996;75(3):177-182.
135. Chan DY, Chan CC, Au DK. Motor relearning programme for stroke patients: a randomized, controlled trial. *Clin Rehabil*. 2006;20(3):191-200.
136. Chan SC, Chan AP. Rehabilitation outcomes following traumatic spinal cord injury in a tertiary spinal cord injury centre: a comparison with an international standard. *Spinal Cord*. 2005;43(8):489-498.
137. Chang WC, Chan C. Rasch analysis for outcomes measures: some methodological considerations. *Archives of Physical Medicine and Rehabilitation*. 1995;76(10):934-9.

138. Chang WC, Slaughter S, Cartwright D, Chan C. Evaluating the FONE FIM: Part I. Construct validity. *Journal of Outcome Measurement*. 1997;1(3):192-218.
139. Chang WC, Slaughter S, Cartwright D, Chan C. Evaluating the FONE FIM: Part II. Concurrent validity & influencing factors. *Journal of Outcome Measurement*. 1997;1(4):259-285.
140. Charpentier P, Legall M, Bernot R. [Comparative study of the Functional Independence Measure and of the Coubert Scale, using the international classification of handicaps]. *Annales de Readaptation et de Médecine Physique*. 1994;37(1):61-67.
141. Chau N, Daler S, Andre JM, Patris A. Inter-rater agreement of two functional independence scales: the Functional Independence Measure (FIM) and a subjective uniform continuous scale. *Disability and Rehabilitation*. 1994;16(2):63-71.
142. Chen CC, Heinemann AW, Granger CV, Linn RT. Functional gains and therapy intensity during subacute rehabilitation: a study of 20 facilities. *Arch Phys Med Rehabil*. Nov 2002;83(11):1514-1523.
143. Chen CL, Tang FT, Chen HC, Chung CY, Wong MK. Brain lesion size and location: effects on motor recovery and functional outcome in stroke patients. *Arch Phys Med Rehabil*. 2000; 81(4):447-452.
144. Chen IC, Cheng PT, Hu AL, et al. Balance evaluation in hemiplegic stroke patients. *ChanGung Med J*. 2000;23(6):339-347.
145. Cheng SK, Man DW. Management of impaired self-awareness in persons with traumatic brain injury. *Brain Inj*. 2006;20(6):621-628.
146. Cherkassky T, Oksenberg A, Froom P, Ring H. Sleep-related breathing disorder and rehabilitation outcome of stroke patients: a prospective study. *Am J Phys Med Rehabil*. 2003;82(6):452-455.
147. Cherney LR, Halper AS, Kwasnica CM, Hervey RL, Zhang M. Recovery of functional status after right hemisphere stroke: relationship with unilateral neglect. *Arch Phys Med Rehabil*. 2001;82(3):322-328.
148. Childers MK, Brashear A, Jozefczyk P, et al. Dose-dependent response to intramuscular botulinum toxin type A for upper-limb spasticity in patients after a stroke. *Arch Phys Med Rehabil*. 2004;85(7):1063-1069.
149. Cho DY, Chen CC, Chang CS, Lee WY, Tso M. Endoscopic surgery for spontaneous basal ganglia hemorrhage: comparing endoscopic surgery, stereotactic aspiration, and craniotomy in noncomatose patients. *Surg Neurol*. 2006;65(6):547-555.
150. Choo B, Umraw N, Gomez M, Cartotto R, Fish JS. The utility of the functional independence measure (FIM) in discharge planning for burn patients. *Burns*. 2006;32(1):20-23.
151. Cifu DX, Huang ME, Kolakowsky-Hayner SA, Seel RT. Age, outcome, and rehabilitation costs after paraplegia caused by traumatic injury of the thoracic spinal cord, conus medullaris, and cauda equina. *J Neurotrauma*. 1999;16(9):805-815.

152. Cifu DX, Kreutzer JS, Kolakowsky-Hayner SA, Marwitz JH, Englender J. The relationship between therapy intensity and rehabilitative outcomes after traumatic brain injury: a multicenter analysis. *Arch Phys Med Rehabil.* 2003;84(10):1441-1448.
153. Cifu DX, Seel RT, Kreutzer JS, McKinley WO. A multicenter investigation of age-related differences in lengths of stay, hospitalization charges, and outcomes for a matched tetraplegia sample. *Arch Phys Med Rehabil.* 1999;80(7):733-740.
154. Cisternino MD, Giaquinto S, Maiolo I, Palma E, Valeriani M, Vittoria E. On the outcome in stroke patients one year later: the role of atrial fibrillation. *Eur J Neurol.* 2003;10(1):67-70.
155. Claesson L, Svensson E. Measures of order consistency between paired ordinal data: application to the Functional Independence Measure and Sunnaas index of ADL. *J Rehabil Med.* 2001;33(3):137-144.
156. Clair AA, O'Konski M. The effect of rhythmic auditory stimulation (RAS) on gait characteristics of cadence, velocity, and stride length in persons with late stage dementia. *J Music Ther.* 2006;43(2):154-163.
157. Cohen ME, Marino RJ. The tools of disability outcomes research functional status measures. *Arch Phys Med Rehabil.* Dec 2000;81(12 Suppl 2):S21-29.
158. Cook A, Hussey S. Delivering assistive technology services to the consumer . In: Cook A, Hussey S., eds. *Assistive Technologies: Principles and Practice*. St. Louis: C. V. Mosby, 1995: pp.141-202.
159. Cook L, Smith DS, Truman G. Using Functional Independence Measure profiles as an index of outcome in the rehabilitation of brain-injured patients. *Archives of Physical Medicine and Rehabilitation.* 1994;75(4):390-393.
160. Cooke DM, McKenna K, Fleming J, Darnell R. Construct and ecological validity of the occupational therapy adult perceptual screening test (OT-APST). *Scan J Occup Ther.* 2006;13(1):49-61.
161. Copes WS, Stark MM, Lawnick MM et al. Linking data from national trauma and rehabilitation registries. *Journal of Trauma.* 1996;40(3):428-436.
162. Copes WS, Staz CF, Konvolinka CW, Sacco WJ. American College of Surgeons audit filters: associations with patient outcome and resource utilization. *Journal of Trauma.* 1995;38(3):432-438.
163. Corrigan JD, Smith-Knapp K, Granger CV. Validity of the functional independence measure for persons with traumatic brain injury. *Archives of Physical Medicine and Rehabilitation.* 1997;78(8):828-34.
164. Corrigan JD, Smith-Knapp K, Granger CV. Outcomes in the first 5 years after traumatic brain injury. *Arch Phys Med Rehabil.* 1998;79(3):298-305.
165. Costa BA, Engrav LH, Holavanahalli R, et al. Impairment after burns: a two-center, prospective report. *Burns.* 2003;29(7):671-675.

166. Coster WJ, Haley SM, Jette AM. Measuring patient-reported outcomes after discharge from inpatient rehabilitation settings. *J Rehabil Med.* 2006;38(4):237-242.
167. Coster WJ, Haley SM, Ludlow LH, Andres PL, Ni PS. Development of an applied cognition scale to measure rehabilitation outcomes. *Arch Phys Med Rehabil.* 2004;85(12):2030-2035.
168. Cotter EM, Burgio LD, Stevens AB, Roth DL, Gitlin LN. Correspondence of the functional independence measure (FIM) self-care subscale with real-time observations of dementia patients' ADL performance in the home. *Clin Rehabil.* 2002;16(1):36-45.
169. Cowen TD, Huang CT, Lebow J, DeVivo MJ, Hawkins LN. Functional outcomes after inpatient rehabilitation of patients with end-stage renal disease. *Archives of Physical Medicine and Rehabilitation.* 1995;76(4):355-359.
170. Cowen TD, Meythaler JM, DeVivo MJ, Ivie CS 3rd., Lebow J, Novack TA. Influence of early variables in traumatic brain injury on Functional Independence Measure scores and rehabilitation length of stay and charges. *Archives of Physical Medicine and Rehabilitation.* 1995;76(9):797-803.
171. Crewe NM, Dijkers M. Functional assessment. In: Cushman LA, Scherer MJ, editors. *Psychological Assessment in Medical Rehabilitation. Measurement and Instrumentation in Psychology.* American Psychological Association, Washington, DC, XV, 1995: pp. 101-144.
172. Cromwell SJ, Paquette VL. The effect of botulinum toxin A on the function of a person with poststroke quadriplegia. *Physical Therapy.* 1996;76(4):395-402.
173. Cruse JM, Lewis RE, Bishop GR, Kliesch WF, Gaitan E. Neuroendocrine-immune interactions associated with loss and restoration of immune system function in spinal cord injury and stroke patients. *Immunologic Research.* 1992;11(2):104-116.
174. Cruse JM, Lewis RE Jr., Bishop GR, Kliesch WF, Gaitan E, Britt R. Decreased immune reactivity and neuroendocrine alterations related to chronic stress in spinal cord injury and stroke patients. *Pathobiology.* 1993;61(3-4):183-192.
175. Cully JA, Gfeller JD, Heise RA, Ross MJ, Teal CR, Kunik ME. Geriatric depression, medical diagnosis, and functional recovery during acute rehabilitation. *Arch Phys Med Rehabil.* 2005;86(12):2256-2260.
176. Cushman LA, Scherer MJ. Measuring the relationship of assistive technology use, functional status over time, and consumer-therapist perceptions of ATs. *Assist Technol.* 1996;8(2):103-109.
177. Cusick CP, Brooks CA, Whiteneck GG. The use of proxies in community integration research. *Arch Phys Med Rehabil.* 2001;82(8):1018-1024.
178. Cusick CP, Gerhart KA, Mellick DC. Participant-proxy reliability in traumatic brain injury outcome research. *J Head Trauma Rehabil.* 2000;15(1):739-749.

179. Czyrny JJ, Hamilton BB, Gresham GE. Rehabilitation of the stroke patient. *Advances in Clinical Rehabilitation*. 1990;64-87.
180. Czyrny JJ, Merrill A. Rehabilitation of amputees with end-stage renal disease: functional outcome and cost. *American Journal of Physical Medicine and Rehabilitation*. 1994;73(5):353-357.
181. Dahmer ER, Shilling MA, Hamilton BB, Bontke CF, Englander J, Kreutzer JS, Ragnarsson KT, Rosenthal M. A model systems database for traumatic brain injury. *Journal of Head Trauma Rehabilitation*. 1993;8(2):12-25.
182. D'Alisa S, Baudo S, Mauro A, Miscio G. How does stroke restrict participation in long-term post-stroke survivors? *Acta Neurol Scand*. 2005;112(3):157-162.
183. Dallmeijer AJ, Dekker J, Roorda LD, et al. Differential item functioning of the Functional Independence Measure in higher performing neurological patients. *J Rehabil Med*. 2005;37(6):346-352.
184. D'Aduila MA, Smith T, Organ D, Lichtman S, Reding M. Validation of a lateropulsion scale for patients recovering from stroke. *Clin Rehabil*. 2004;18(1):102-109.
185. Dario A, Scamoni C, Bono G, Ghezzi A, Zaffaroni M. Functional improvement in patients with severe spinal spasticity treated with chronic intrathecal baclofen infusion. *Funct Neurol*. 2001;16(4):311-315.
186. Daverat P, Petit H, Kemoun G, Dartigues JF, Barat M. The long term outcome of 149 patients with spinal cord injury. *Paraplegia*. 1995;33:665-668.
187. Davidoff GN, Roth EJ, Haughton JS, Ardner MS. Cognitive dysfunction in spinal cord injury patients: sensitivity of the Functional Independence Measure subscales vs. neuropsychologic assessment. *Archives of Physical Medicine and Rehabilitation*. 1990;71(5):326-329.
188. Davidoff GN, Roth EJ, Richards S. Cognitive deficits in spinal cord injury: epidemiology and outcome. *Archives of Physical Medicine and Rehabilitation*. 1992;73(3):275-284.
189. de Langen EG, Frommelt P, Wiedmann KD, Amann J. Evaluation of functional independence in rehabilitation by the Functional Independence Measure (FIM). *Rehabilitation*. 1995;34(1):IV-XI.
190. Daving Y, Andren E, Nordholm L, Grimby G. Reliability of an interview approach to the Functional Independence Measure. *Clin Rehabil*. 2001;15(3):301-310.
191. Davis CH, Fardanesh L, Rubner D, Wanlass RL, McDonald CM. Profiles of functional recovery in fifty traumatically brain-injured patients after acute rehabilitation. *Am J Phys Med Rehabil*. 1997;76(3):213-218.
192. Deathe B, Miller WC, Speechley M. The status of outcome measurement in amputee rehabilitation in Canada. *Arch Phys Med Rehabil*. 2002;83(7):912-918.

193. de Guise E, Leblanc J, Feyz M, Lamoureaux J. Prediction of the level of cognitive functional independence in acute care following traumatic brain injury. *Brain Inj.* 2005;19(13):1087-1093.
194. Dellarosa DM, Chan RSK, Toglia JP, Finkelstein N. ADL assessment in acute care: simultaneous grading of physical and verbal levels of assistance. *Occupational Therapy Practice.* 1991;2(2):38-45.
195. DeJong G, Horn SD, Smout RJ, Ryser DK. The early impact of the inpatient rehabilitation facility prospective payment system on stroke rehabilitation case mix, practice patterns, and outcomes. *Arch Phys Med Rehabil.* 2005;86(12 Suppl 2):s93-s100.
196. DeJong G, Hsieh CH, Putman K, Smout RJ, Horn SD, Tian W. Physical therapy activities in stroke, knee arthroplasty, and traumatic brain injury rehabilitation: their variation, similarities, and association with functional outcomes. *Phys Ther.* 2011;91(12):1826-37.
197. de Langen EG, Frommelt P. [Evaluation of functional independence in rehabilitation by the Functional Independence Measure (FIM)]. *Die Rehabilitation.* 1995;34(1):4-11.
198. Demers L, Giroux F. Validity of the Functional Independence Measure (FIM) for follow up of rehabilitation in the elderly. *Canadian Journal on Aging.* 1997;16(4):626-646.
199. Deng AW, Wei D, Zhang JH, Ran CF, Wang M. [Rehabilitation therapy in early stage following spinal cord injuries]. *Di Yi Jun Yi Da Xue Xue Bao.* 2004;24(6):706-707,710.
200. Deng AW, Yuang XG, Wei D, Zhang JH, Ran CF, Wang M. [Effect of power-frequency electromagnetic fields on stroke during rehabilitation]. *Di Yi Jun Yi Da Xue Xue Bao.* 2004;24(8):946-949,952.
201. Denys P, Azouvi P, Denormandie P, Samuel C, Patel A, Bussel B. Late cognitive and behavioral improvement following treatment of disabling orthopaedic complications of a severe closed head injury. *Brain Inj.* 1996;10(2):149-153.
202. de Seze M, Wiart L, Bon-Saint-Come A, et al. Rehabilitation of postural disturbances of hemiplegic patients by using trunk control retraining during exploratory exercises. *Arch Phys Med Rehabil.* 2001;82(6):793-800.
203. Deshpande SA, MacNeill SE, Lichtenberg PA, Pithadia J, Velez L. Functional outcome differences in acute versus subacute geriatric rehabilitation. *Topics in Geriatric Rehabilitation.* 1998; 13(4):30-38.
204. Desrosiers J, Bravo G, Hebert R, Dutil E, Mercier L. Validation of the Box and Block Test as a measure of dexterity of elderly people: reliability, validity, and norms studies. *Archives of Physical Medicine & Rehabilitation.* 1994;75(7):751-755.
205. Desrosiers J, Rochette A, Noreau L, Bravo G, Hebert R, Boutin C. Comparison of two functional independence scales with a participation measure in post-stroke rehabilitation. *Arch Gerontol Geriatr.* 2003;37(2):157-172.

206. Deutsch A, Fiedler RC, Iwanenko W, Granger CV, Russell CF. The Uniform Data System for Medical Rehabilitation report: patients discharged from subacute rehabilitation programs in 1999. *Am J Phys Med Rehabil.* Sep 2003;82(9):703-711.
207. Deutsch A, Braun S, Granger CV. The Functional Independence Measure (FIM<sub>SM</sub> Instrument) and the Functional Independence Measure for Children (WeeFIM Instrument): ten years of development. *Archives of Physical Medicine and Rehabilitation.* 1996; 77(12):1226-32.
208. DeVivo MJ, Richards JS, Storer SL, Go BK. Spinal cord injury: rehabilitation adds life to years. *The Western Journal of Medicine.* 1991;154(5):602-606.
209. Diamond PT, Felsenthal G, Macciocchi SN, Butler DH, Lally-Cassady D. Effect of cognitive impairment on rehabilitation outcome. *American Journal of Physical Medicine and Rehabilitation.* 1996;75(1):40-43.
210. Diamond PT, Holroyd S, Macciocchi SN, Felsenthal G. Prevalence of depression and outcome on the geriatric rehabilitation unit. *American Journal of Physical Medicine and Rehabilitation.* 1995;74(3):214-217.
211. Diamond PT, Conaway MR, Mody SH, Bhirangi K. Influence of hemoglobin levels on inpatient rehabilitation outcomes after total knee arthroplasty. *J Arthroplasty.* 2006;21(5):636-641.
212. Diard C, Ravaud JF, Held JP. French survey of postpolio sequelae. Risk factors study and medical social outcome. *American Journal of Physical Medicine and Rehabilitation.* 1994;73(4):264-267.
213. DiBlasi M, Savage J. Revitalizing a documentation system. *Rehabilitation Nursing.* 1992;17(1):27-29.
214. Dickson HG, Hodgkinson A, Kohler F. Inpatient quality assurance by local analysis of uniform data set data. *Journal of Quality in Clinical Practice.* 1994;14(3):145-148.
215. Dickson HG, Kohler F. The Functional Independence Measure: a comparative validity and reliability study [letter; comment]. *Disability and Rehabilitation.* 1995;17(8):456.
216. Dickson HG, Kohler F. The multi-dimensionality of the FIM motor items precludes an interval scaling using Rasch analysis. *Scandinavian Journal of Rehabilitation Medicine.* 1996;28(3):159-162.
217. Dickson HG, Kohler F. The multi-dimensionality of the FIM motor items precludes an interval scaling using Rasch analysis [comment]. *Scandinavian Journal of Rehabilitation Medicine.* 1997;29(4):267-72.
218. Dickson HG, Kohler F. Inter-rater reliability of the 7-level Functional Independence Measure (FIM). *Scandinavian Journal of Rehabilitation Medicine.* 1995;27(4):253-4.
219. Dickson HG, Kohler F. Functional independence measure (FIM). *Scand J Rehabil Med.* 1999;31(1):63-64.



220. Dijkers MP. A computer adaptive testing simulation applied to the FIM instrument motor component. *Arch Phys Med Rehabil.* 2003;84(3):384-393.
221. Dijkers MP, Yavuzer G. Short versions of the telephone motor Functional Independence Measure for use with persons with spinal cord injury. *Arch Phys Med Rehabil.* 1999;80(11):1477-1484.
222. Dijkers MP, Yavuzer G, Ergin S, Weitzenkamp D, Whiteneck GG. A tale of two countries; environmental impacts on social participation after spinal cord injury. *Spinal Cord.* 2002;40(7):351-362.
223. Diringer MN, Edwards DF, Zazulia AR. Hydrocephalus: a previously unrecognized predictor of poor outcome from supratentorial intracerebral hemorrhage. *Stroke.* 1998;29(7):1352-1357.
224. Disler PB, Roy CW, Smith BP. Predicting hours of care needed. *Archives of Physical Medicine and Rehabilitation.* 1993;74(2):139-143.
225. Ditunno JF. Functional assessment measures in CNS trauma. *Journal of Neurotrauma.* 1992;9(Suppl 1):S301-S305.
226. Ditunno JF, Cohen ME, Formal CH, Whiteneck GG. Functional outcomes. In: Stover SL, DeLisa JA, Whiteneck GG, eds. *Spinal Cord Injury Clinical Outcomes from the Model Systems.* Gaithersburg, MD: Aspen Publishers. 1995:170-184.
227. Ditunno JF, Ditunno PL, Graziani V, et al. Walking index for spinal cord injury (WISCI): an international multicenter validity and reliability study. *Spinal Cord.* 2000;38(4):234-243.
228. Ditunno JF, Marino RJ, Crozier KS. Neurological and functional assessments in acute spinal injury: uses in prognosis and management. *Trauma Quarterly.* 1993;9(2) 44-52.
229. Ditunno JF, Young W, Donovan W, Creasy G. The international standards booklet for neurological and functional classification of spinal cord injury. *Paraplegia.* 1994;32(2):70-80.
230. Dobkin B, Apple D, Barbeau H, et al,. Weight-supported treadmill vs over-ground training for walking after acute incomplete SCI. *Neurology.* 2006;66(4):484-493.
231. Dobrez DG, Lo Sasso AT, Heinemann AW. The effect of prospective payment on rehabilitative care. *Arch Phys Med Rehabil.* 2004;85(12):1909-1914.
232. Doctor JN, Wolfson AM, McKnight P, Burns SP. The effect of inaccurate FIM instrument ratings on prospective payment: a study of clinician expertise and FIM rating difficulty as contributing to inaccuracy. *Arch Phys Med Rehabil.* 2003;84(1):46-50.
233. Dodd KJ, Morris ME. Lateral pelvic displacement during gait: abnormalities after stroke and changes during the first month of rehabilitation. *Arch Phys Med Rehabil.* 2003;84(8):1200-1205.

234. Dodds TA, Martin DP, Stolov WC, Deyo RA. A validation of the Functional Independence Measurement and its performance among rehabilitation inpatients. *Archives of Physical Medicine and Rehabilitation*. 1993;74(5):531-536.
235. Dombovy ML, Drew-Cates J, Serdars R. Recovery and rehabilitation following subarachnoid haemorrhage: Part II. Long-term follow-up. *Brain Inj*. 1998;12(10):887-894.
236. Dombovy ML, Olek AC. Recovery and rehabilitation following traumatic brain injury. *Brain Inj*. 1997;11(5):305-318.
237. Donaghy S, Wass PJ. Interrater reliability of the Functional Assessment Measure in a brain injury rehabilitation program. *Arch Phys Med Rehabil*. 1998;79(10):1231-1236.
238. Donnelly C, Eng JJ, Hall J, et al. Client-centered assessment and the identification of meaningful treatment goals for individuals with a spinal cord injury. *Spinal Cord*. 2004;42(5):302-307.
239. Drew Cates J. Functional Assessment In: Stanley M, Beare PG, ed. *Gerontological Nursing*. Philadelphia: F. A. Davis, 1995:91-97.
240. Dromerick AW, Edwards DF, Diringer MN. Sensitivity to changes in disability after stroke: a comparison of four scales useful in clinical trials. *J Rehab Res Dev*. 2003;40(1):1-8.
241. Duarte E, Marco, Muniesa JM, et al. Trunk control test as a function predictor in stroke patients. *J Rehabil Med*. 2002;34(6):267-272.
242. Dubljanin-Raspopović E, Marković-Denić L, Nikolić D, et al. Is anemia at admission related to short-term outcomes of elderly hip fracture patients? *Cent Eur J Med*. 2011;6(4):483-489.
243. Duncan PW. Stroke disability. *Physical Therapy*. 1994;74(5):399-407.
244. Duncan PW, Reker DM, Horner RD, et al. Performance of a mail-administered version of a stroke-specific outcome measure, the Stroke Impact Scale. *Clin Rehabil*. 2002;16(5):493-505.
245. Dunkerley AL, Ashburn A, Stack EL. Deltoid triceps transfer and function independence of people with tetraplegia. *Spinal Cord*. 2000;38(7):435-441.
246. Duong TT, Englander J, Wright J, Cifu DX, Greenwald BD, Brown AW. Relationship between strength, balance, and swallowing deficits and outcome after traumatic brain injury: a multicenter analysis. *Arch Phys Med Rehabil*. 2004;85(8):1291-1297.
247. Duran FS, Lugo L, Ramirez L, Eusse E. Effects of an exercise program on the rehabilitation of patients with spinal cord injury. *Arch Phys Med Rehabil*. 2001;82(10):1349-1354.
248. Eastwood EA, Hagglund KJ, Ragnarsson KT, Gordon WA, Marino RJ. Medical rehabilitation length of stay and outcomes for persons with traumatic spinal cord injury – 1990-1997. *Arch Phys Med Rehabil*. 1999;80(11):1457-1463.

249. Edwards DF, Chen YW, Diringer MN. Unified Neurological Stroke Scale is valid in ischemic and hemorrhagic stroke. *Stroke*. 1995;26(10):1852-1858.
250. Eilersten TB, Kramer AM, Schlenker RE, Hrinkevich CA. Application of functional independence measure-function related groups and resource utilization groups-version III systems across post-acute settings. *Med Care*. 1998;36(5):695-705.
251. Eisermann U, Haase I, Klandy B. Computer-aided multimedia training in orthopedic rehabilitation. *Am J Phys Med Rehabil*. 2004;83(9):670-680.
252. Eldar R, Ring H, Tshuwa M, Dynia A, Ronan R. Quality of care for urinary incontinence in a rehabilitation setting for patients with stroke. Simultaneous monitoring of process and outcome. *Int J Qual Health Care*. 2001;13(1):57-61.
253. Emhoff TA, McCarthy M, Cushman M, Garb JL, Valenziano C. Functional scoring of multi-trauma patients: who ends up where? *Journal of Trauma*. 1991;31(9):1227-1232.
254. Englander J, Cifu DX, Wright JM, Black K. The association of early computed tomography scan findings and ambulation, self-care, and supervision needs at rehabilitation discharge and at 1 year after traumatic brain injury. *Arch Phys Med Rehabil*. 2003;84(2):214-220.
255. Ergeletzis D, Kevorkian CG, Rintala D. Rehabilitation of the older stroke patient: functional outcome and comparison with younger patients. *Am J Phys Med Rehabil*. 2002;81(12):881-889.
256. Essengen S, Seckin U, Borman P, Bodur H, Kutsal YG. Drug consumption in a group of elderly residents of a nursing home: relationship to cognitive impairment and disability. *J Am Med Dir Assoc*. 2000;1(5):197-201.
257. Evans LK, Yurkow J, Siegler EL. The CARE Program: a nurse-managed collaborative outpatient program to improve function of frail older people. Collaborative Assessment and Rehabilitation for Elders. *Journal of the American Geriatrics Society*. 1995;43(10):1155-1160.
258. Falconer JA, Naughton BJ, Dunlop DD, Roth EJ, Strasser DC, Sinacore JM. Predicting stroke inpatient rehabilitation outcome using a classification tree approach. *Archives of Physical Medicine and Rehabilitation*. 1994;75(6):619-625.
259. Feld JA, Rabadi MH, Blau AD, Jordan BD. Berg balance scale and outcome measures in acquired brain injury. *Neurorehabil Neural Repair*. 2001;15(3):239-244.
260. Ferguson KA, Plando G, Kobetic R, Triolo RJ, Marsolais EB. Walking with a hybrid orthosis system. *Spinal Cord*. 1999;37(11):800-804.
261. Ferrio G, Franchignoni F, Benevolo E, Ottonello M, Scocchi M, Xanthi M. The influence of comorbidities and complications on discharge function in stroke rehabilitation inpatients. *Eura Medicophys*. 2006;42(2):91-96.
262. Feys P, Duportail M, Kos D, Van Asch P, Ketelaer P. Validity of the TEMPA for the measurement of upper limb function in multiple sclerosis. *Clin Rehabil*. 2002;16(2):166-173.

263. Fiedler RC, Granger CV. The Functional Independence Measure: a measurement of disability and medical rehabilitation. In: Chino N, Melvin JL, eds. *Functional Evaluation of Stroke Patients*. Tokyo: Springer-Verlag, 1996:75-92.
264. Fiedler RC, Granger CV. Uniform Data System for Medical Rehabilitation: report of first admissions for 1996. *Am J Phys Med Rehabil*. 1998;77(1):69-75.
265. Fiedler RC, Granger CV, Ottenbacher KJ. The Uniform Data System for Medical Rehabilitation: report of first admissions for 1994. *American Journal of Physical Medicine and Rehabilitation*. 1996;75(2):125-129.
266. Fisher AG. The assessment of IADL motor skills: an application of many-faceted Rasch analysis. *American Journal of Occupational Therapy*. 1993;47(4):319-329.
267. Fisher CG, Noonan VK, Smith DE, Wing PC, Dvorak MF, Kwon B. Motor recovery, functional status, and health-related quality of life in patients with complete spinal cord injuries. *Spine*. 2005;30(19):220-2207.
268. Fisher WP, Harvey RF, Taylor P, Kilgore KM, Kelly CK. Rehabits: a common language of functional assessment. *Archives of Physical Medicine and Rehabilitation*. 1995;76(2):113-122.
269. Flanders AE, Spettell CM, Friedman DP, Marino RJ, Herbson GJ. The relationship between the functional abilities of patients with cervical spinal cord injury and the severity of damage revealed by MR imaging. *Am J Neuroradiol*. 1999;20(5):926-934.
270. Foley MA, Shahrokhi K, Robinson KM. A behavioral neuropsychological approach. Strategies for the difficult geriatric patient. *Topics in Geriatric Rehabilitation*. 1990;5(3):32-42.
271. Fong KN, Chan CC, Au DK. Relationship of motor and cognitive abilities to functional performance in stroke rehabilitation. *Brain Inj*. 2001;15(5):443-453.
272. Forer S. How to make program evaluation work for you: utilization for program service management. *Neuro Rehabilitation*. 1992;2:52-71.
273. Forrest G, Nagao M, Iqbal A, Kakar R. In patient rehabilitation of patients requiring hemodialysis: improving efficiency of care. *Arch Phys Med Rehabil*. 2005;86(10):1949-1952.
274. Forrest GP. Inpatient rehabilitation of patients requiring hemodialysis. *Arch Phys Med Rehabil*. 2004;85(1):51-53.
275. Fortis P, Maravita A, Gallucci M, et al. Rehabilitating patients with left spatial neglect by prism exposure during a visuomotor activity. *Neuropsychology*. 2010;24(6):681-97.
276. Foxx-Orenstein A, Kolakowsky-Hayner S, Marwitz JH, et al. Incidence, risk factors, and outcomes of fecal incontinence after acute brain injury; findings from the Traumatic Brain Injury Model Systems national database. *Arch Phys Med Rehabil*. 2003;84(2):231-237.

277. Franchignoni FP, Benevolo E, Zelaschi GP, DiPatrizi S, Sessarego S, Tesio L. Recovery from physical disability as evaluated by the Functional Independence Measure inpatient stroke rehabilitation. *Europa Medicophysica*. 1995;31(2):67-75.
278. Franchignoni FP, Tesio L, Ricupero C, Martino MT. Trunk Control Test as an early predictor of stroke rehabilitation outcomes. *Stroke*. 1997;28(7):1328-1385.
279. Franchignoni F, Tesio L, Martino MT, Benevolo E, Castagna M. Length of stay of stroke rehabilitation inpatients: prediction through the functional independence measure. *Ann 1<sup>st</sup> Super Sanita*. 1998;34(4):463-467.
280. Franchignoni F, Ottonello M, Benevolo E, Tesio L. Satisfaction with hospital rehabilitation: is it related to life satisfaction, functional status, age, or education? *J Rehabil Med*. 2002;34(3):105-108.
281. Franchignoni F, Tesio L, Benevolo E, Ottonello M. Psychometric properties of the Rivermead Mobility Index in Italian stroke rehabilitation inpatients. *Clin Rehabil*. 2003;17(3):273-82.
282. Franchignoni F, Orlandini D, Ferriero G, Moscato TA. Reliability, validity, and responsiveness of the locomotor capabilities index in adults with lower-limb amputation undergoing prosthetic training. *Arch Phys Med Rehabil*. 2004;85(5):743-748.
283. Francisco G, Chae J, Chawla H, et al. Electromyogram-triggered neuromuscular stimulation for improving the arm function of acute stroke survivors: a randomized plot study. *Arch Phys Med Rehabil*. 1998;79(5):570-575.
284. Frankel JE, Marwitz JH, Cifu DX, Kreutzer JS, Englander J, Rosenthal M. A follow-up study of older adults with traumatic brain injury: taking into account decreasing length of stay. *Arch Phys Med Rehabil*. 2006;87(1):57-62.
285. Franzoi AC, Araujo AQ. Disability profile of patients with HTLV-I-associated myelopathy/tropical spastic paraparesis using Functional Independence Measure (FIM). *Spinal Cord*. 2005;43(4):236-240.
286. Frattali CM. Functional assessment of communication: merging public policy with clinical views. *Aphasiology*. 1992;6(1):63-83.
287. Frattali CM, Chapey R, Sacchett C, Marshall J, Scherzer E, Worrall L. Clinical forum: functional assessment of communication: merging public policy with clinical views. *Aphasiology*. 1992;6(1):63-83,85-93,95-103,105-116.
288. Frattali CM. Perspectives on functional assessment and its use for policy making. *Disability and Rehabilitation*. 1993;15(1):1-9.
289. Freeman JA, Playford ED, Nicholas RS, Thompson AJ. A neurological rehabilitation unit: audit of activity and outcome. *Journal of the Royal College of Physicians of London*. 1996;30(1):21-26.
290. Freeman JA, Hobart JC, Playford ED, Undy B, Thompson AJ. Evaluating neurorehabilitation: lessons from routine data collection. *J Neurol Neurosurg Psychiatry*. 2005;76(5):723-728.

291. Fricke J. Measuring outcomes in rehabilitation: a review. *British Journal of Occupational Therapy*. 1993;56(5):217-221.
292. Fricke J, Unsworth CA. Inter-rater reliability of the original and modified Barthel Index, and a comparison with the Functional Independence Measure. *Australian Occupational Therapy Journal*. 1997;44(1):22-29.
293. Fricke J, Unsworth C, Worrel D. Reliability of the Functional Independence Measure with occupational therapists. *Australian Occupational Therapy Journal*. 1993;40(1):7-15.
294. Frymark TB, Mullen RC. Influence of the prospective payment system on speech-language pathology services. *Am J Phys Med Rehabil*. 2005;84(1):12-21.
295. Fucile J. Functional rehabilitation in cancer care. *Seminars in Oncology Nursing*. 1992;8(3):186-189.
296. Fuhrer MJ. Rehabilitation research in the 1980s. Eisenberg MG, Grzesak RC, eds. *Advances in Clinical Rehabilitation*. New York; Springer Publishing. 1988:232-256.
297. Fuhrer MJ, Garber SL, Rintala DH, Clearman R, Hart KA. Pressure ulcers in community-resident persons with spinal cord injury: prevalence and risk factors. *Archives of Physical Medicine and Rehabilitation*. 1993;74(11):1172-1177.
298. Fuhrer MJ, Rintala DH, Hart KA, Clearman R, Young ME. Depressive symptomatology in persons with spinal cord injury who reside in the community. *Archives of Physical Medicine and Rehabilitation*. 1993;74(3):255-260.
299. Fuhrer MJ, Rintala DH, Hart KA, Clearman R, Young ME. Relationship of life satisfaction to impairment, disability, and handicap among persons with spinal cord injury living in the community. *Archives of Physical Medicine and Rehabilitation*. 1992;73(6):552-557.
300. Fujiwara T, Hara Y, Akaboshi K, Chino N. Relationship between shoulder muscle strength and functional independence measure (FIM) score among C6 tetraplegics. *Spinal Cord*. 1999;37(1):58-61.
301. Gagne DE, Hoppes S. The effects of collaborative goal-focused occupational therapy on self-care skills: a pilot study. *Am J Occup Ther*. 2003;57(2):215-219.
302. Gagnon C, Desrosiers J, Mathieu J. Autosomal recessive spastic ataxia of Charlevoix-Saguenay: upper extremity aptitudes, function independence and social participation. *Int J Rehabil Res*. 2004;27(3):253-256.
303. Gagnon C, Mathieu J, Desrosiers J. Standardized finger-nose test validity for coordination assessment in an ataxic disorder. *Can J Neurol Sci*. 2004;31(4):484-489.
304. Gagnon D, Nadeau S, Tom V. Clinical and administrative outcomes during publicly-funded inpatient stroke rehabilitation based on a case-mix group classification model. *J Rehabil Med*. 2005;37(1):45-52.

305. Galin LC, Rode G, Soler Michel P, Eyssette M, Boisson D. Interest in the study of the Functional Independence Measure (FIM) in the evaluation of the rehabilitation for hemiplegic patients over the age of 60. *Annales De Readaptation Et De Medecine Physique*. 1994;37(5):281-290.
306. Galloway RV, Granger CV, Karmarkar AM, et al. The Uniform Data System for Medical Rehabilitation: Report of patients with debility discharged from inpatient rehabilitation programs in 2000-2010. *Am J Phys Med Rehabil*. 2013;92:14-27.
307. Gallien P, Sebert E, Galanth E, Brissot R, Perrigot M. Relevance of Functional Independence Measurement in taking care of incontinence in stroke. *Annales De Readaptation Et De Medecine Physique*. 1995;38(1):21-24.
308. Gallien P, Aghulon C, Durufle A, et al. Magnetoencephalography in stroke: a 1-year follow-up study. *Eur J Neurol*. 2003;10(4):373-382.
309. Galynker I, Cohen L, Salvit C, et al. Psychiatric symptom severity and length of stay on an intensive rehabilitation unit. *Psychosomatics*. 2000;41(2):114-120.
310. Garcia RA, Gaebler-Spira D, Sisung C, Heinemann AW. Functional improvement after pediatric spinal cord injury. *Am J Phys Med Rehabil*. 2002;81(6):458-463.
311. Gauggel S, Bocker M, Zimmerman P, Privou C, Lutz D. [Item response theory and its application in neurology. Measurement of activity limitations in neurologic patients]. *Nervenarzt*. 2004;75(12):1179-1186.
312. Gennarelli TA, Champion HR, Copes WS, Sacco WJ. Comparison of mortality, morbidity, and severity of 59,713 head injured patients with 114,447 patients with extracranial injuries. *Journal of Trauma*. 1994;7(6):962-968.
313. George ER, Scholten DJ, Buechler CM, Jordan-Tibbs J, Mattice C, Albrecht RM. Failure of methylprednisolone to improve the outcome of spinal cord injuries. *American Surgeon*. 1995;61(8):659-663 (discussion 663-664).
314. Gerrard P, Goldstein R, **DiVita MA**, Ryan CM, **Mix J**, **Niewczyk P**, Kazis L, Kowalske K, Schneider JC, Zafonte R. Validity and reliability of the FIM instrument in the inpatient burn population. *Am J Phys Med Rehabil*. 2013;Mar 5 epub ahead of print.
315. Ghislandi S, Apolone G, Garattini L, Ghislandi I. Is EQ-5D a valid measure of HRQoL in patients with movement disorders? A comparison with SF-36 and FIM questionnaires. *Eur J Health Econ*. 2002;3(2):125-130.
316. Gialanella B, Benvenuti P, Santoro R, Scolari S. [Disability and rehabilitation outcome in hemiplegia with global aphasia and neglect]. *Clin Ter*. 2003;154(4):237-243.
317. Gialanella B, Monguzzi V, Santoro R, Rocchi S. Functional recovery after hemiplegia in patients with neglect: the rehabilitative role of anosognosia. *Stroke*. 2005;35(12):2687-2690.
318. Giaquinto S. Comorbidity in post-stroke rehabilitation. *Eur J Neurol*. 2003;10(3):235-238.

319. Giaquinto S. Death or improvement: the fate of highly disabled patients after stroke rehabilitation. *Clin Exp Hypertens*. 2006;28(3-4):357-364.
320. Giaquinto S, Buzzelli S, DiFrancesco L, et al. On the prognosis of outcome after stroke. *Acta Neurol Scand*. 1999;100(3):202-208.
321. Giaquinto S, Buzzelli S, DiFrancesco L, Villani S. Recovery mechanisms are stronger than expected. *Arch Gerontol Geriatr Suppl*. 2004;9:217-222.
322. Giaquinto S, Ferrara I, Muschera R, Pagano G, Nolfe G. The effects of atrial fibrillation on functional recovery in post-stroke patients. *Disabil Rehabil*. 2001;23(5):204-208.
323. Giaquinto S, Fraioli L. Enhancement of the somatosensory N140 components during attentional training after stroke. *Clin Neurophysiol*. 2003;114(2):329-335.
324. Giquinto S, Majolo II, Palma E, Roncacci S, Sciarra A, Vittoria E. Very old people can have favorable outcome after hip fracture: 58 patients referred to rehabilitation. *Arch Gerontol Geriatr*. 2000;31(1):13-18.
325. Giaquinto S, Mascio M, DiLibero F, Fagnoli M, Pittiglio S. Is the Kennard effect supported by clinical evidence? *Arch Gerontol Geriatr Suppl*. 2004;9:223-228.
326. Giaquinto S, Mascio M, Fraioli L. The physiopathological bases of recovery processes: the bases of stroke rehabilitation. The CASSINO project. Computer assisted somatosensory stimulation inducing new organizations. *Clin Exp Hypertens*. 2002;24(7-8):543-553.
327. Gillen R, Tennen H, McKee T. Unilateral spatial neglect: relation to rehabilitation outcomes in patients with right hemisphere stroke. *Arch Phys Med Rehabil*. 2005;86(4):763-767.
328. Gillen R, Tennen H, McKee TE, Gernert-Dott P, Affleck G. Depressive symptoms and history of depression predict rehabilitation efficiency in stroke patients. *Arch Phys Med Rehabil*. 2001;82(12):1645-1649.
329. Glass CA, Tesio L, Itzkovich M, et al. Spinal Cord Independence Measure, version III: applicability to the UK spinal cord injured population. *J Rehabil Med*. 2009;41(9):723-8.
330. Glass RM. Program evaluation. In: England B, Glass RM, Patterson CH, eds. *Quality Rehabilitation: Results-Oriented Patient Care*. Chicago: American Hospital Association, 1989:19-37.
331. Gokkaya NK, Aras MD, Cacki A. Health-related quality of life of Turkish stroke survivors. *Int J Rehabil Res*. 2005;28(3):229-235.
332. Goldstein FC, Strasser DC, Woodard JL, Roberts VJ. Functional outcome of cognitively impaired hip fracture patients on a geriatric rehabilitation unit. *J Am Geriatr Soc*. 1997;45(1):35-42.
333. Goldstein TS. Measuring Function. In: *Functional Rehabilitation on Orthopedics*. Gaithersburg, MD: Aspen Publishers. 1995:183-215.



334. Gonnella C. Program evaluation. In: Fletcher GF, Banja JD, Jann BB, Wolf SL, eds. *Rehabilitation Medicine: Contemporary Clinical Perspectives*. Philadelphia, PA: Lea and Febiger, 1992:243-268.
335. Gosman-Hedstrom G, Blomstrand C. Evaluation of a 5-level functional independence measure in a longitudinal study of elderly stroke survivors. *Disabil Rehabil*. 2004;26(7):410-418.
336. Gosman-Hedstrom G, Svensson E. Parallel reliability of the functional independence measure and the Barthel ADL index. *Disabil Rehabil*. 2000;22(16):702-715.
337. Gowland C, Stratford P, Ward M, Moreland J, Torresin W, Van Hullenar S, Sanford J, Barreca S, Vanspall B, Plews N. Measuring physical impairment and disability with the Chedoke-McMaster Stroke Assessment. *Stroke*. 1993;24(1):58-63.
338. Grace J, Nadler JD, White DA, et al. Folstein vs. modified Mini-Mental State Examination in geriatric stroke: stability, validity, and screening utility. *Archives of Neurology*. 1995;52(5):477-484.
339. Grade C, Redford B, Chrostowski J, Toussaint L, Blackwell B. Methylphenidate in early poststroke recovery: a double-blind, placebo-controlled study. *Arch Phys Med Rehabil*. 1998;79(9):1047-1050.
340. Graham PH, Capp A, Delaney G, et al. A pilot randomized comparison of dexamethasone 96mg vs 16 mg per day for malignant spinal-cord compression treated by radiotherapy: TROG 01.05 Superdex study. *Clin Oncol (R Coll Radiol)*. 2006;18(1):70-76.
341. Granger CV. Health accounting - functional assessment of the long term patient. In: Kotte FJ, ed. *Krusen's Handbook of Physical Medicine and Rehabilitation 4th ed*. Philadelphia: W B Saunders. 1990:270-28.
342. Granger CV. Rehabilitation of the aged. In: Calkins E, Ford AB, Katz PR, eds. *Practice of Geriatrics Second Edition*. Philadelphia: W B Saunders Co. 1992:162-170.
343. Granger CV. The emerging science of functional assessment: our tool for outcomes analysis. *Archives of Physical Medicine and Rehabilitation*. 1998;79(3):298-305.
344. Granger CV, Braun SL. Outcome assessments and measures in pediatrics. In: Betz, RR, ed. *The Child with a Spinal Cord Injury*. Rosemond, IL: American Academy of Orthopaedic Surgeons. 1996:709-718.
345. Granger CV, Brownschidle CM. Outcome measurement in medical rehabilitation. *International Journal of Technology Assessment in Health Care*. 1995;11(2):262-268.
346. Granger CV, Clark GS. Functional status and outcomes of stroke rehabilitation. *Topics in Geriatric Rehabilitation*. 1994;9(3):72-84.
347. Granger CV, Cotter AC, Hamilton BB, Fiedler RC. Functional assessment scales: a study of persons after stroke. *Archives of Physical Medicine and Rehabilitation*. 1993;74(2):133-138.

348. Granger CV, Cotter AC, Hamilton BB, Fielder RC, Hens MM. Functional assessment scales: a study of persons with multiple sclerosis. *Archives of Physical Medicine and Rehabilitation*. 1990;71(11):870-875.
349. Granger CV, Deutsch A, Linn RT. Rasch analysis of the Functional Independence Measure (FIM) Mastery Test. *Archives of Physical Medicine and Rehabilitation*. 1998;79(1):52-57.
350. Granger CV, Deutsch A, Tesio L. [Uniform Data System]. *Il Fisioterapista*. 1998;4(5):1-9.
351. Granger CV, Divan N, Fiedler RC. Functional assessment scales. A study of persons after traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 1995;74(2):107-113.
352. Granger CV, Gresham GE. Functional assessment in rehabilitation medicine. *Physical Medicine and Rehabilitation Clinics of North America*. 1993;4(3):417-423.
353. Granger CV, Hamilton BB. Measurement of stroke rehabilitation outcome in the 1980s. *Stroke*. 1990;21(Suppl II):II-46-II-47.
354. Granger CV, Hamilton BB. The Uniform Data System for Medical Rehabilitation report of first admissions for 1991. *American Journal of Physical Medicine and Rehabilitation*. 1993;72(1):33-38.
355. Granger CV, Hamilton BB. The Uniform Data System for Medical Rehabilitation report of first admissions for 1990. *American Journal of Physical Medicine and Rehabilitation*. 1992;71(2):108-113.
356. Granger CV, Hamilton BB. The Uniform Data System for Medical Rehabilitation report of first admissions for 1992. *American Journal of Physical Medicine and Rehabilitation*. 1994;73(1):51-55.
357. Granger CV, Hamilton BB, Fielder RC. Discharge outcome after stroke rehabilitation. *Stroke*. 1992;23(7):978-982.
358. Granger CV, Hamilton BB, Keith RA, Zielezny M, Sherwin FS. Advances in functional assessment for medical rehabilitation. *Topics in Geriatric Rehabilitation*. 1986;1(3):59-74.
359. Granger CV, Hamilton BB, Linacre JM, Heinemann AW, Wright BD. Performance profiles of the Functional Independence Measure. *American Journal of Physical Medicine and Rehabilitation*. 1993;72(2):84-89.
360. Granger CV, Hens MM, Brownschidle CM. Disability and rehabilitation. In: Halbreich U, ed. *Multiple Sclerosis: A Neuropsychiatric Disorder*. Washington DC: American Psychiatric Press. 1993:137-161.
361. Granger CV, Kelly-Hayes M, Johnston M, Deutsch A, Braun S, Fiedler RC. Quality and outcome measures in medical rehabilitation. In: Braddom RL, ed. *Physical Medicine and Rehabilitation*. Philadelphia: W. B. Saunders. 1996:239-253.

362. Granger CV, Mann WC, Ottenbacher KJ, Tomita MR, Fiedler RC. Functional measures of geriatric subjects in the community. *Topics in Geriatric Rehabilitation*. 1994;10(1):7-21.
363. Granger CV, Ottenbacher KJ, Baker JG, Sehgal A. Reliability of a brief outpatient functional outcomes assessment measure. *American Journal of Physical Medicine and Rehabilitation*. 1994;74(6):469-475.
364. Granger CV, Ottenbacher KJ, Fiedler RC. The Uniform Data System for Medical Rehabilitation report of first admissions for 1993. *American Journal of Physical Medicine and Rehabilitation*. 1995;74(1):62-66.
365. Granger CV, Wende K, Brownscheidle CM. Use of the FIM instrument in a trial of intramuscular interferon beta-1a for disease progression in relapsing-remitting multiple sclerosis. *Am J Phys Med Rehabil*. Jun 2003;82(6):427-436.
366. Granger CV, Karmarkar AM, Graham JE, et al. The Uniform Data System for Medical Rehabilitation: Report of patients with traumatic spinal cord injury discharged from rehabilitation programs in 2002-2010. *Am J Phys Med Rehabil*. 2012;91:289-299.
367. Gray DS, Burnham RS. Preliminary outcome analysis of a long-term rehabilitation program for severe acquired brain injury. *Arch Phys Med Rehabil*. 2000;81(11):1447-1456.
368. Greenberg E, Treger I, Ring H. Rehabilitation outcomes in patients with brain tumors and acute stroke: comparative study of inpatient rehabilitation. *Am J Phys Med Rehabil*. 2006;85(7):568-673.
369. Green D, Lawler M, Rosen MS, et al. Recombinant human erythropoietin: effect on the functional performance of anemic orthopedic patients. *Archives of Physical Medicine and Rehabilitation*. 1996;77(3):242-246.
370. Greenspan AI, Wrigley JM, Kresnow M, Branche-Dorsey CM, Fine PR. Factors influencing failure to return to work due to traumatic brain injury. *Brain Injury*. 1996;10(3):207-218.
371. Greenwald BD, Seel RT, Cifu DX, Shah AN. Gender-related differences in acute rehabilitation lengths of stay, charges, and functional outcomes for a matched sample with spinal cord injury: a multicenter investigation. *Arch Phys Med Rehabil*. 2001;82(9):1181-1187.
372. Greve KW, Lindberg RF, Bianchini KJ, Adams D. Construct validity and predictive value of the Hooper Visual Organization Test in stroke rehabilitation. *Appl Neuropsychol*. 2000;7(4):215-222.
373. Grey N, Kennedy P. The Functional independence Measure: a comparative study of clinician and self ratings. *Paraplegia*. 1993;31(7):457-461.
374. Grigsby J, Kooken R, Hershberger J. Simulated neural networks to predict outcomes, costs, and length of stay among orthopedic rehabilitation patients. *Archives of Physical Medicine and Rehabilitation*. 1994;75(10):1077-1081.

375. Grill E, Stucki G, Scheuringer M, Melvin J. Validation of International Classification of Functioning, Disability, and Health (ICF) Core Sets for early postacute rehabilitation facilities: comparisons with three other functional measures. *Am J Phys Med Rehabil*. 2006;85(8):640-649.
376. Grimby G, Andren E, Daving Y, Wright B. Dependence and perceived difficulty in daily activities in community-living stroke survivors 2 years after stroke: a study of instrumental structures. *Stroke*. 1998; 29(9):1843-1849.
377. Grimby G, Andren E, Holmgren E, Wright B, Linacre JM, Sundh V. Structure of a combination of Functional Independence Measure and Instrumental Activity Measure items in community-living persons: a study of individuals with cerebral palsy and spina bifida. *Archives of Physical Medicine and Rehabilitation*. 1996;77(11):1109-1114.
378. Grimby G, Gudjonsson G, Rodhe M, Sunnerhagen KS, Sundh V, Ostensson ML. The Functional Independence Measure in Sweden: experience for outcome measure in rehabilitation medicine. *Scandinavian Journal of Rehabilitation Medicine*. 1996;28(2):51-62.
379. Grisson SP, Dunagen L. Improved satisfaction during inpatient rehabilitation after hip and knee arthroplasty: a retrospective analysis. *Am J Phys Med Rehabil*. 2001;80(11):798-803.
380. Gross JC. A comparison of the characteristics of incontinent and continent stroke patients in a rehabilitation program. *Rehabil Nurs*. 1998;23(3):132-140.
381. Gross JC. Urinary incontinence and stroke outcomes. *Arch Phys Med Rehabil*. 2000;81(1):22-27.
382. Gross JC, Faulkner EA, Goodrich SW, Kain ME. A patient acuity and staffing tool for stroke rehabilitation inpatients based on the FIM instrument. *Rehabilitation Nursing*. 2001;26(3):108-13.
383. Gross JC, Goodrich SW, Kain ME, Faulkner EA. Determining stroke rehabilitation inpatients' level of nursing care. *Clin Nurs Res*. 2001;10(1):40-51.
384. Guccione AA. Functional assessment of the elderly. In: Guccione AA, ed. *Geriatric Physical Therapy*. St. Louis: C. V. Mosby, 1993:113-123.
385. Guerini Rocco D, Mercieri A, Yavuzer G. Multidimensional health-status assessment of chronic hemodialysis patients: the impact on quality of life. *Eura Medicophys*. 2006;42(2):113-119.
386. Guillaume D, Van Havenbergh A, Vloeberghs M, Vidal J, Roeste G. A clinical study of intrathecal baclofen using a programmable pump for intractable spasticity. *Arch Phys Med Rehabil*. 2005;86(11):2165-2171.
387. Guo Y, Collaco CR, Bruera E. Heterotopic ossification in critical illness and cancer: a report of 2 cases. *Arch Phys Med Rehabil*. 2002;83(6):855-859.

388. Gurka JA, Felmingham KL, Baguley IJ, Schotte DE, Crooks J, Marssozeky JE. Utility of the functional assessment measure after discharge from inpatient rehabilitation. *J Head Trauma Rehabil.* 1999;14(3):247-256.
389. Haas U, Mayer H, Evers CG. [Interobserver reliability of the 'Functional Independence Measure' (FIM) in patients with craniocerebral injuries]. *Pflege.* 2002;15(4):191-197.
390. Hachisuka K, Okazaki T, Ogata H. Self-rating Barthel Index compatible with the original Barthel Index and the Functional Independence Measure motor score. *Journal of UOEH Association of Health Sciences.* 1997;19(2):107-121.
391. Hachisuka K, Tsutsui Y, Furusawa K, Ogata H. Gender differences in disability and lifestyle among community-dwelling elderly stroke patients in Kitakyushu, Japan. *Arch Phys Med Rehabil.* 1998;79(8):992-1002.
392. Haley SM, Coster WJ, Binda-Sundberg K. Measuring physical disablement: the contextual challenge. *Physical Therapy.* 1994;74(5):443-451.
393. Haley SM, Coster WJ, Ludlow LH. Pediatric functional outcome measures. *Physical Medicine and Rehabilitation Clinics of North America.* 1991;2(4):689-723.
394. Hall KM. Overview of functional assessment scales in brain injury rehabilitation. *NeuroRehabilitation.* 1992;2(4):98-113.
395. Hall KM, Hamilton BB, Gordon WA, Zalsler ND. Characteristics and comparisons of functional assessment indices: Disability Rating Scale, Functional Independence Measure, and Functional Assessment Measure. *Journal of Head Trauma Rehabilitation.* 1993;8(2):60-74.
396. Hall KH, Johnson MV. Outcomes evaluation in TBI rehabilitation - part II: measurement tools for a nationwide data system. *Archives of Physical Medicine and Rehabilitation.* 1994;75(suppl.):SC10-SC18.
397. Hall KM, Mann N, High WM, Wright J, Kreutzer JS, Wood D. Functional measures after traumatic brain injury: ceiling effects of FIM, FIM+FAM, DRS, and CIQ. *Journal of Head Trauma Rehabilitation.* 1996;11(5):27-39.
398. Hall KM, Cohen ME, Wright J, Call M, Werner P. Characteristics of the Functional Independence Measure in traumatic spinal cord injury. *Arch Phys Med Rehabil.* 1999;80(11):1471-1476.
399. Hall KM, Bushnik T, Lakisic-Kazazic B, Wright J, Cantagallo A. Assessing traumatic brain injury outcome measures for long-term follow-up of community-based individuals. *Arch Phys Med Rehabil.* 2001;82(3):367-374.
400. Hall KM, Wallbom AS, Englander J. Premorbid history and traumatic brain injury. *Neurorehabilitation.* 1998;10(1):3-12.
401. Halper AS, Cherney LR, Cichowski K, Zhang M. Dysphagia after head trauma: the effect of cognitive-communicative impairments on functional outcomes. *J Head Trauma Rehabil.* 1999;14(5):486-496.

402. Hamilton BB, Deutsch A, Russell C, Fielder RC, Granger CV. Relation of disability costs to function: spinal cord injury. *Archives of Physical Medicine and Rehabilitation*. 1999;80(4):385-91.
403. Hamilton BB, Laughlin JA, Fielder RC, Granger CV. Inter-rater reliability of the 7-level Functional Independence Measure (FIM). *Scandinavian Journal of Rehabilitation Medicine*. 1994;26(3):115-119.
404. Hamilton BB, Granger CV. Totaled functional score can be valid [letter; comment]. *Archives of Physical Medicine and Rehabilitation*. 1989;70(12):861-863.
405. Hamilton BB, Granger CV. Disability outcomes following inpatient rehabilitation for stroke. *Physical Therapy*. 1994;74(5):494-503.
406. Hamilton BB, Granger CV, Sherwin FS, Zielezny M, Tashman JS. Uniform national data system for medical rehabilitation. In: Fuhrer MJ, ed. *Rehabilitation Outcomes: Analysis and Measurement*. Baltimore, MD: Paul H. Brookes Publishing Co., 1987:137-147.
407. Hamilton BB, Laughlin JA, Fiedler RC, Granger CV. Interrater Reliability of the 7-level Functional Independence Measure (FIM). *Scandinavian Journal of Rehabilitation Medicine*. 1994;26(3):115-119.
408. Hammond FM, Hart T, Busnik T, Corrigan JD, Sasser H. Change and predictors of change in communication, cognition, and social function between 1 and 5 years after traumatic brain injury. *J Head Trauma Rehabil*. 2004;19(4):314-328.
409. Han L, Law-Gibson D, Reding M. Key neurological impairments influence function-related group outcomes after stroke. *Stroke*. 2002;33(7):1920-1924.
410. Hanger HC, Whitewood P, Brown G, et al. A randomized controlled trial of strapping to prevent post-stroke shoulder pain. *Clin Rehabil*. 2000;14(4):370-380.
411. Harada N, Kominski G, Sofaer S. Development of a resource-based patient classification scheme for rehabilitation. *Inquiry*. 1993;30:54-63.
412. Hart T, Millis S, Novack T, Englander J, Fidler-Sheppard R, Bell KR. The relationship between neuropsychologic function and levels of caregiver supervision at 1 year after traumatic brain injury. *Arch Phys Med Rehabil*. 2003;84(2):221-230.
413. Hartman-Maeir A, Soroker N, Oman SD, Katz N. Awareness of disabilities in stroke rehabilitation—a clinical trial. *Disabil Rehabil*. 2003;25(1):35-44.
414. Harvey RL, Roth EJ, Heineman AW, Lovell LL, McGuire JR, Diaz S. Stroke rehabilitation: clinical predictors of resource utilization. *Arch Phys Med Rehabil*. 1998;79(11):1349-1355.
415. Hashimoto K, Okamoto T, Watanabe S, Ohashi M. Effectiveness of a comprehensive day treatment program for rehabilitation of patients with acquired brain injury in Japan. *J Rehabil Med*. 2006;38(1):20-25.

416. Hastings J, Gowans S, Watson DE. Effectiveness of occupational therapy following organ transplantation. *Can J Occup Ther*. 2004;71(4):238-242.
417. Hawley CA. Return to driving after head injury. *J Neurol Neurosurg Psychiatry*. 2001;70(6):761-766.
418. Hecox R, Roach KE, DasVarma JM, Giraud JE, Davis CM, Neulen K. Functional Independence Measurement (FIM) of patients receiving Easy Street - a retrospective study. *Physical and Occupational Therapy in Geriatrics*. 1994;12(3):17-31.
419. Heinemann AW, Hamilton BB, Linacre JM, Wright BD, Granger CV. Functional status and therapeutic intensity during inpatient rehabilitation. *American Journal of Physical Medicine and Rehabilitation*. 1995;74(4):315-326.
420. Heinemann AW, Kan E. Relationship between cognitive impairment and disability following stroke and traumatic brain injury. *Europa Medicophysica*. 1996;32(4):195-205.
421. Heinemann AW, Kirk P, Hastie BA, et al. Relationships between disability measures and nursing effort during medical rehabilitation for patients with traumatic brain and spinal cord injury. *Archives of Physical Medicine and Rehabilitation*. 1997;78(2):143-149.
422. Heinemann AW, Linacre JM, Wright BD, Hamilton BB, Granger CV. Measurement characteristics of the Functional Independence Measure. *Topics in Stroke Rehabilitation*. 1994;1(3):1-15.
423. Heinemann AW, Linacre JM, Wright BD, Hamilton BB, Granger CV. Prediction of rehabilitation outcomes with disability measures. *Archives of Physical Medicine and Rehabilitation*. 1994;75(2):133-143.
424. Heinemann AW, Linacre JM, Wright BD, Hamilton BB, Granger CV. Relationships between impairment and physical disability as measured by the Functional Independence Measure. *Archives of Physical Medicine and Rehabilitation*. 1993;74(6):566-573.
425. Hens MM. Functional evaluation. In: Dittmar SS, ed. *Rehabilitation Nursing: Process and Application*. St. Louis: C. V. Mosby Co. 1989:486-501.
426. Herbold JA, Bonistall K, Walsh MB. Rehabilitation following total knee replacement, total hip replacement, and hip fracture: a case controlled comparison. *J Geriatr Phys Ther*. 2011;34(4):155-60.
427. Heriseanu R, Baguley IJ, Slewa-Younan S. Two-point discrimination following traumatic brain injury. *J Clin Neurosci*. 2005;12(2):156-160.
428. Herrman N, Black SE, Lawrence J, Szekely C, Szalai JP. The Sunnybrook Stroke Study: a prospective study of depressive symptoms and functional outcome. *Stroke*. 1998;29(3):618-624.
429. HersHKovitz A, Beloosesky Y, Brill S, Gottlieb D. Is a day hospital rehabilitation programme associated with reduction of handicap in stroke patients? *Clin Rehabil*. 2004;18(3):261-266.

430. Herskovitz A, Gottlieb D, Beloosesky Y, Brill S. Programme evaluation of a geriatric rehabilitation day hospital. *Clin Rehabil.* 2003;17(7):750-755.
431. Herskovitz A, Pulatov I, Brill S, Beloosesky Y. Can hip-fractured elderly patients maintain their rehabilitation achievements after 1 year? *Disabil Rehabil.* 2012;34(4):304-10.
432. Heruti RJ, Lusky A, Barell V, Ohry A, Adunsky A. Cognitive status at admission: does it affect the rehabilitation outcome of elderly patients with hip fracture? *Arch Phys Med Rehabil.* 1999;80(4):432-436.
433. Heruti RJ, Lusky A, Danker R, et al. Rehabilitation outcome of elderly patients after a first stroke: effect of cognitive status at admission on the functional outcome. *Arch Phys Med Rehabil.* 2002;83(6):742-749.
434. Hetherington H, Earlam RJ. Measurement of disability after multiple injuries: The Functional Independence Measure. *European Journal of Surgery, Acta Chirurgica.* 1995;161(8):549-55.
435. Hetherington H, Earlam RJ. Rehabilitation after injury and the need for coordination. *Injury.* 1994;25(8):527-31.
436. Hetherington H, Earlam RJ, Kirk CJ. The disability status of injured patients measured by the Functional Independence Measure (FIM) and their use of rehabilitation services. *Injury.* 1995;26(2):97-101.
437. Hiller S, English C, Crotty M, Segal L, Berhardt J, Esterman A. Circuit class or seven-day therapy for increasing intensity of rehabilitation after stroke: protocol of the CIRCUIT trial. *Int J of Stroke.* 2011;6(6):560-565.
438. Hinkle JL, Forbes E. Pilot project on functional outcome in stroke. *Journal of Neuroscience Nursing.* 1996;28(1):13-18.
439. Hinkle JL. A descriptive study of function in acute motor stroke. *West J Nurs Res.* 2001;23(3):296-312.
440. Hinkle JL. Variables explaining functional recovery following motor stroke. *J Neurosci Nurs.* 2006;38(1):6-12.
441. Hinkle JL, Davies J, Ng D, McClaran J. Examining assessment tools for discharge planning. *Nurs Times.* 2008;104(43):32-5.
442. Hobart JC, Lamping DL, Freeman JA, et al. Evidence-based measurement: which disability scale for neurologic rehabilitation? *Neurology.* 2001;57(4):639-644.
443. Hobart JC, Cano SJ, Thompson AJ. Effect sizes can be misleading: is it time to change the way we measure change? *J Neurol Neurosurg Psychiatry.* 2010;81:1044-1048.
444. Hoenig H, McIntyre L, Sloan R, Branch LG, Truncali A, Horner RD. The reliability of a self-reported measure of disease, impairment, and function in persons with spinal cord dysfunction. *Archives of Physical Medicine and Rehabilitation.* 1998;79(4):378-387.



445. Hoenig H, Duncan PW, Horner RD, et al. Structure, process, and outcomes in stroke rehabilitation. *Med Care*. 2002;40(11):1036-1047.
446. Holland A, O'Conner RJ, Thompson AJ, Playford ED, Hobart JC. Talking the talk on walking the walk: A 12-item generic walking scale suitable for neurological conditions? *J Neurol*. 2006;253(12):1594-1602.
447. Hoppe H, Reifshauer A, Conradi E. Functional Independence Measure - a new assessment to evaluate the success of rehabilitation. *Physikalische Medizin Rehabilitationsmedizin Kurortmedizin*. 1994;4(4):128-130.
448. Houlden H, Edwards M, McNeil J, Greenwood R. use of the Barthel index and the Functional Independence Measure during early inpatient rehabilitation after single incident brain injury. *Clin Rehabil*. 2006;20(2):153-159.
449. Hsueh IP, Lin JH, Jeng JS, Hsieh CL. Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke. *J Neurol Neurosurg Psychiatry*. 2002;73(2):188-190.
450. Huang ME, Cifu DX, Keyser-Marcus L. Functional outcome after brain tumor and acute stroke: a comparative analysis. *Arch Phys Med Rehabil*. 1998;79(11):1386-1390.
451. Huang ME, Cifu DX, Keyser-Marcus L. Functional outcomes in patients with brain tumor after inpatient rehabilitation: comparison with traumatic brain injury. *Am J Phys Med Rehabil*. 2000;79(4):327-335.
452. Huang ME, Johns JS, White J, Sanford K. Venous thromboembolism in a rehabilitation setting after major lower-extremity amputation. *Arch Phys Med Rehabil*. 2005;86(1):73-78.
453. Huang ME, Wartalla JE, Kreutzer JS. Functional outcomes and quality of life in patients with brain tumors: a preliminary report. *Arch Phys Med Rehabil*. 2001;82(11):1540-1546.
454. Hubbard W, Bauer D. Programme evaluation: what we can learn from the United States experience. *Australian Journal of Physiotherapy*. 1993;39(1):53-57.
455. Huebner RA, Johnson K, Bennett CM, Schneck C. Community participation and quality of life outcomes after adult traumatic brain injury. *Am J Occup Ther*. 2003;57(2):177-185.
456. Hurvitz EA, Conti GE, Brown SH. Changes in movement characteristics of the spastic upper extremity after botulinum toxin injection. *Arch Phys Med Rehabil*. 2003;84(3):444-454.
457. Ickenstein GW, Stein J, Ambrosi D, Goldstein R, Horn M, Bogdahn U. Predictors of survival after severe dysphagic stroke. *J Neurol*. 2005;252(12):1510-1516.
458. Iizuka R, Reding M. Use of percutaneous endoscopic gastrostomy feeding tubes and functional recovery in stroke rehabilitation: a case-matched controlled study. *Arch Phys Med Rehabil*. 2005;86(5):1049-1052.

459. Inoye M, Hashimoto H, Mio T, Sumin K. Influence of initial status on functional gain for Japanese patients with first cerebral hemorrhage. *J Rehabil Med*. 2001;33(1):12-15.
460. Inoye M, Hashimoto H, Mio T, Sumino K. Influence of admission functional status on functional change after stroke rehabilitation. *Am J Phys Med Rehabil*. 2001;80(2):121-125; quiz 126, 146.
461. Inoye M, Kishi K, Ikeda Y, et al. Prediction of functional outcome after stroke rehabilitation. *Am J Phys Med Rehabil*. 2000;79(6):513-518.
462. Itzkovich M, Tripolski M, Zeilig G, et al. Rasch analysis of the Catz-Itzkovich spinal cord independence measure. *Spinal Cord*. 2002;40(8):396-407.
463. Ivie CS, De Vivo MJ. Predicting unplanned hospitalizations in persons with spinal cord injury. *Archives of Physical Medicine and Rehabilitation*. 1994;75(11):1182-1188.
464. Jackson JP, Schkade JK. Occupational Adaptation model versus biomechanical rehabilitation model in the treatment of patients with hip fractures. *Am J Occup Ther*. 2001;55(5):531-537.
465. James R, Gines D, Menlove A, Horn SD, Gassaway J, Smout RJ. Nutrition support (tube feeding) as a rehabilitation intervention. *Arch Phys Med Rehabil*. 2005;86(12suppl2):s82-s92.
466. Jansen DA, Cimprich B. Attentional impairment in persons with multiple sclerosis. *Journal of Neuroscience Nursing*. 1994;26(2):95-102.
467. Jaworski D, Kult T, Boynton PR. The Functional Independence Measure: a pilot study comparison of observed and reported ratings. *Rehabilitation Nursing Research*. 1994;3(4):141-147.
468. Jensen AP, Abresch RT, Carter GT. The reliability and validity of a self-report version of the FIM instrument in persons with neuromuscular disease and chronic pain. *Arch Phys Med Rehabil*. 2005;86(1):116-122.
469. Jensen MP, Abresch RT, Carter GT. The reliability and validity of a self-report version of the FIM instrument in persons with neuromuscular disease and chronic pain. *Arch Phys Med Rehabil*. 2005;86(1):116-122.
470. Jette DU, Warren RL, Wirtalla C. The relation between therapy intensity and outcomes of rehabilitation in skilled nursing facilities. *Arch Phys Med Rehabil*. 2005;86(3):373-379.
471. Jette DU, Warren RL, Wirtalla C. Functional independence domains in patients receiving rehabilitation in skilled nursing facilities: evaluation of psychometric properties. *Arch Phys Med Rehabil*. 2005;86(6):1089-94.
472. Johns JS, Cifu DX, Keyser-Marcus L, Jolles PR, Fratkin MJ. Impact of clinically significant heterotopic ossification on functional outcome after traumatic brain injury. *J Head Trauma Rehabil*. 1999;14(3):269-276.

473. Johnston MV, Findley TW, DeLuca J, Katz RT. Research in physical medicine and rehabilitation: XII. Measurement tools with application to brain injury. *American Journal of Physical Medicine and Rehabilitation*. 1991;70(1):40-56.
474. Johnston MV, Granger CV. Outcomes research in medical rehabilitation: a primer and introduction to a series. *American Journal of Physical Medicine and Rehabilitation*. 1994;73(4):296-303.
475. Johnston MV, Hall KM. Outcomes evaluation in TBI rehabilitation -- part I: overview and system principles. *Archives of Physical Medicine and Rehabilitation*. 1994;75(Suppl):SC2-SC9.
476. Johnston MV, Keith RA. Measurement standards for medical rehabilitation and applications. *Physical Medicine and Rehabilitation Clinics of North America*. 1993;4(3):425-449.
477. Johnston MV, Keith RA, Hinderer S. Measurement standards for interdisciplinary medical rehabilitation. *Archives of Physical Medicine and Rehabilitation*. 1992;73(Suppl):S3-S23.
478. Johnston MV, Kirshblum S, Zorowitz R, Schiflett SC. Prediction of outcomes following rehabilitation of stroke patients. *NeuroRehabilitation*. 1992;2:72-97.
479. Johnston MV, Wilkerson DL, Maney M. Evaluation of the quality and outcome of medical rehabilitation programs. In: DeLisa J, Gans BM, Currie DM, eds. *Rehabilitation Medicine: Principles and Practice*. Philadelphia: J B Lippincott, 1993:240-268.
480. Johnston MV, Wood KD, Fielder R. Characteristics of effective and efficient rehabilitation programs. *Arch Phys Med Rehabil*. 2003;84(3):410-418.
481. Johnston MV, Wood K, Millis S, Page S, Chen D. Perceived quality of care and outcomes following spinal cord injury: minority status in the context of multiple predictors. *J Spinal Cord Med*. 2004;27(3):241-251.
482. Johnston MV, Shawaryn MA, Malec J, Kreutzer J, Hammond FM. The structure of functional and community outcomes following traumatic brain injury. *Brain Inj*. 2006;20(4):391-407.
483. Jones CA, Feeny DH. Agreement between patient and proxy responses during recovery after hip fracture: evidence for the FIM Instrument. *Archives of Physical Medicine and Rehabilitation*. 2006;87(10):1382-1387.
484. Jones GR, Miller TA, Petrella RJ. Evaluation of rehabilitation outcomes in older patients with hip fractures. *Am J Phys Med Rehabil*. 2002;81(7):489-497.
485. Jong CN, Zafonte RD, Millis SR, Yavuzer G. The effect of cocaine on traumatic brain injury outcome: a preliminary evaluation. *Brain Inj*. 1999;12(12):1017-1023.
486. Jongjit J, Komsopapong L, Songjakkaew P, Kongsakon R. Health-related quality of life after hip fracture in the elderly community-dwelling. *Southeast Asian J Trop Med Public Health*. 2003;34(3):670-674.

487. Juneja G, Czrny JJ, Linn RT. Admission balance and outcomes of patients admitted for acute inpatient rehabilitation. *Am J Phys Med Rehabil*. 1998;77(5):388-393.
488. Kamenetz Y, Beloosesky Y, Zeltzer C, et al. Relationship between routine hematological parameters, serum IL-3, IL-6 and erythropoietin and mild anemia and degree of function in the elderly. *Aging (Milano)*. 1998;10(1):32-38.
489. Kamikubo T, Ohashi M, Hashimoto K, Miyano S. [Cognitive dysfunction in 32 diffuse axonal injury cases]. *No To Shinkei*. 2003;55(8):669-673.
490. Kamikubo T, Ohashi M, Hashimoto K, Okamoto T, Miyano S. [Lowered VIQ level following traumatic brain injury]. *No To Shinkei*. 2004;56(11):952-956.
491. Kapidzic-Durakovic S, Zonic-Imamovic M, Mulaosmanovic A. Family participation in determination of FIM and Barthel indices. *Bosn J Basic Med Sci*. 2006;6(2):51-53.
492. Kaplan CP, Corrigan JD. The relationship between cognition and functional independence in adults with traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 1994;75(6):643-647.
493. Karamahmetoglu SS, Karacan I, Elbasi N, Demirel G, Koyuncu H, Dosoglu M. The functional independence measure in spinal cord injured patients: comparison of questioning with observational rating. *Spinal Cord*. 1997;35(1):22-25.
494. Karatas M, Cetin N, Bayramoglu M, Delik A. Trunk muscle strength in relation to balance and functional disability in unihemispheric stroke patients. *Am J Phys Med Rehabil*. 2004;83(2):81-87.
495. Karatas M, Dilek A, Erkan H, Yavuz N, Sozay S, Akman N. Functional outcome in stroke patients with atrial fibrillation. *Arch Phys Med Rehabil*. 2000;81(8):1025-1029.
496. Kaszniak A, Bortz J. Issues in evaluating the cost-effectiveness of neuropsychological assessment in rehabilitation settings. In: Glueckauf R, Secrest L, Bond G, McDonel E, eds. *Improving Assessment in Rehabilitation and Health*. Newbury Park, CA: Sage Publications, 1993:176-211.
497. Katz DI, White DK, Alexander MP, Klein BB. *Recovery of ambulation after traumatic brain injury*. *Arch Phys Med Rehabil*. 2004;85(6):865-869.
498. Keith R. Comprehensive rehabilitation: themes, models, and issues. In: Eisenberg M, Glueckauf R, Zaretsky H, eds. *Medical Aspects of Disability: A Handbook for the Rehabilitation Professional*. New York: Springer Publishing Co., 1993:3-22.
499. Keith RA, Granger CV, Hamilton BB, Sherwin FS. The Functional Independence Measure: a new tool for rehabilitation. *Advances in Clinical Rehabilitation*. 1987;1:6-18.
500. Keith RA, Wilson DB, Gutierrez P. Acute and subacute rehabilitation for stroke: a comparison. *Archives of Physical Medicine and Rehabilitation*. 1995;76:495-500.
501. Kelley FA, Kawamoto TT, Rubenstein LZ. Assessment of the geriatric patient. In: Kiernat JM, ed. *Occupational Therapy and the Older Adult: A Clinical Manual*. Gaithersburg, MD: Aspen Publishers, 1991:76-98.

502. Kelly PJ, Furie KL, Shafqat S, Rallis N, Chang Y, Stein J. Functional recovery following rehabilitation after hemorrhagic and ischemic stroke. *Arch Phys Med Rehabil.* 2003;84(7):968-972.
503. Kelly PJ, Stein J, Shafqat S, et al. Functional recovery after rehabilitation for cerebellar stroke. *Stroke.* 2001;32(2):530-534.
504. Kelly S, Jessop EG. A comparison of measures of disability and health status in people with physical disabilities undergoing vocational rehabilitation. *J Public Health Med.* 1996;18(2):169-174.
505. Kelly-Hayes M. Functional Evaluation. In: Hoeman SP, ed. *Rehabilitation Nursing: Process and Application (2nd ed.)*. St. Louis: C. V. Mosby Co. 1996:144 -155.
506. Kennedy RE, Livingston L, Marwitz JH, Gueck S, Kreutzer JS, Sander AM. Complicated mild traumatic brain injury on the inpatient rehabilitation unit: a multicenter analysis. *J Head Trauma Rehabil.* 2006;21(3):260-271.
507. Kersten P, McLellan L, George S, Smith JA. The Southampton Needs Assessment Questionnaire (SNAQ): a valid tool for assessing the rehabilitation needs of the disabled people. *Clin Rehabil.* 2000;14(6):641-650.
508. Kesiktas N, Paker N, Erdogan N, Gulsen G, Bicki D, Yilmaz H. The use of hydrotherapy for the management of spasticity. *Neurorehabil Neural Repair.* 2004;18(4):268-273.
509. Kevorkian CG, Ergelezis D, Rintala D. Nonagenarians on a rehabilitation unit: characteristics, progress, and outcomes. *Am J Phys Med Rehabil.* 2004;83(4):266-272.
510. Keyser-Marcus LA, Bricout JC, Wehman P, et al. Acute predictors of return to employment after traumatic brain injury: a longitudinal follow-up. *Arch Phys Med Rehabil.* 2002;83(5):635-641.
511. Khader MS, Tomlin GS. Change in wheelchair transfer performance during rehabilitation of men with cerebrovascular accident. *American Journal of Occupational Therapy.* 1994;48(10):899-905.
512. Kidd D, Stewart G, Baldry J, et al. The Functional Independence Measure: a comparative validity and reliability study. *Disability and Rehabilitation.* 1995; 17(1):10-14.
513. Kilgore KM, Fisher WP Jr., Silverstein B, Harley JP, Harvey RF. Application of Rasch analysis to the Patient Evaluation and Conference System. *Physical Medicine and Rehabilitation Clinics of North America.* 1993;4(3):493-515.
514. Kiper P, Piron L, Turolla A, Stożek J, Tonin P. The effectiveness of reinforced feedback in virtual environment in the first 12 months after stroke. *Neurol Neurochir Pol.* 2011; 45(5):436-444.
515. Kipnis ND. Accreditation, quality improvement, and program evaluation. In: McCourt AE, ed. *The Specialty Practice of Rehabilitation Nursing: A Core Curriculum*. Skokie IL:

The Rehabilitation Nurses Foundation of the Association of Rehabilitation Nurses. 1993:189-202.

516. Kirby RL, Dupris DJ, Macphee AH, et al. The wheelchair skills test (version 2.4): measurement properties. *Arch Phys Med Rehabil.* 2004;85(5):794-804.
517. Kirk-Sanchez NJ, Roach KE. Relationship between duration of therapy services in a comprehensive rehabilitation program and mobility at discharge in patients with orthopedic problems. *Phys Ther.* 2001;81(3):888-895.
518. Kliesch WF, Cruse JM, Lewis RE, Bishop GR, Brackin B, Lampton JA. Restoration of depressed immune function in spinal cord injury patients receiving rehabilitation therapy. *Paraplegia.* 1996;34(2):82-90.
519. Kosasih JB, Borca HH, Wenninger WJ, Duthie E. Nursing home rehabilitation after acute rehabilitation: predictors and outcomes. *Arch Phys Med Rehabil.* 1998;79(6):670-673.
520. Kovach CR, Puetzer M, Gretzinger P. A descriptive study of nosocomial urinary tract infection in a rehabilitation patient population. *Rehabilitation Nursing Research.* 1993;2(2):81-86.
521. Koyama T, Matsumoto K, Okuno T, Domen K. A new method for predicting functional recovery of stroke patients with hemiplegia: logarithmic modeling. *Clin Rehabil.* 2005;19(7):779-789.
522. Koyama T, Matsumoto K, Okuno T, Domen K. Relationships between independence level of single motor-fim items and fim-motor scores in patients with hemiplegia after stroke: an ordinal logistic modeling study. *J Rehabil Med.* 2006;38(5):280-286.
523. Kucukdeveci AA, Yavuzer G, Elhan AH, Sonel B, Tennant A. Adaptation of the Functional Independence Measure for use in Turkey. *Clin Rehabil.* 2001;15(3):311-319.
524. Kuo YF, Ostir GV, Granger CV, Ottenbacher KJ. Examination of follow-up therapy in patients with stroke. *Am J Phys Med Rehabil.* 2006;85(3):192-200.
525. Kurihara M, Kumagai K, Nakae Y. [Functional outcome after diffuse axonal injury in childhood traumatic brain injury]. *No To Hattatsu.* 1999;31(5):408-414.
526. Kurihara M, Kumagai K, Nakae Y, Noda Y. [Rehabilitation approach to children with sequelae of hypoxic encephalopathy]. *No To Hattatsu.* 1998;30(6):505-511.
527. Kurihara M, Kumagai K, Watanabe M, Noda Y. [Prognosis of severe head injury in childhood: from the viewpoint of brain plasticity]. *No To Hattatsu.* 1996;28(3):243-250.
528. Kwon S, Duncan P, Studentski S, Perera S, Lai SM, Reker D. Measuring stroke impact with SIS: construct validity of SIS telephone administration. *Qual Lif Res.* 2006;15(3):367-376.
529. Kwon S, Hatzema AG, Duncan PW, Min-Lai S. Disability measure in stroke: relationship among the Barthel Index, the Functional Independence Measure, and the Modified Rankin Scale. *Stroke.* 2004;35(4):918-923.

530. Lambert CM, Hurst NP, Lochhead A, McGregor K, Hunter M, Forbes J. A pilot study of the economic cost and clinical outcome of day patient vs inpatient management of active rheumatoid arthritis. *British Journal of Rheumatology*. 1994;33(4):383-388.
531. Lammi MH, Smith VH, Tate RL, Taylor CM. The minimally conscious state and recovery potential: a follow-up study 2 to 5 years after traumatic brain injury. *Arch Phys Med Rehabil*. 2005;86(4):746-754.
532. Landrum PK, Schmidt ND, McLean A, eds. *Outcome-Oriented Rehabilitation: Principles, Strategies, and Tools for Effective Program Management*. Gaithersburg, MD: Aspen Publishers, Inc. 1995.
533. Langdon DW, Thompson AJ. Multiple sclerosis: a preliminary study of selected variables affecting rehabilitation outcome. *Mult Scler*. 1999. 5(2):94-100.
534. Latham NK, Jette DU, Slavin M, et al. Physical therapy during stroke rehabilitation for people with different walking abilities. *Arch Phys Med Rehabil*. 2005;86(12suppl2):s41-s50.
535. Latham NK, Jette DU, Warren RL, Wirtalla C. Pattern of functional change during rehabilitation of patients with hip fracture. *Arch Phys Med Rehabil*. 2006;87(1):111-116.
536. Lawton G, Lundgren-Nilsson A, Biering-Sorensen F, et al. Cross-cultural validity of FIM in spinal cord injury. *Spinal Cord*. 2006;44(12):746-52.
537. Lee L, Goor E, Kennedy C, Walters S, Kirby L. Non-acute case-mix in the Illawarra. *Journal of Quality in Clinical Practice*. 1994;14(1):23-30.
538. Lenze EJ, Munin MC, Dew MA, et al. Adverse effects of depression and cognitive impairment on rehabilitation participation and recovery from hip fracture. *Int J Geriatr Psychiatry*. 2004;19(5):472-478.
539. Lenze EJ, Munin MC, Quear T, et al. Significance of poor patient participation in physical and occupational therapy for functional outcome and length of stay. *Arch Phys Med Rehabil*. 2004;85(10):1599-1601.
540. Lenze EJ, Munin MC, Quear T, et al. The Pittsburg Rehabilitation Participation Scale: reliability and validity of a clinician-related measure of participation in acute rehabilitation. *Arch Phys Med Rehabil*. 2004;85(3):380-384.
541. Leonard H, Fyfe S, Leonard S, Msall M. Functional status, medical impairments, and rehabilitation resources in 84 females with Rett syndrome: a snapshot across the world from the paranatal perspective. *Disabil Rehabil*. 2001;23(3-4):107-117.
542. Leung EC, Rush PJ, Devlin M. Predicting prosthetic rehabilitation outcome in lower limb amputee patients with the functional independence measure. *Arch Phys Med Rehabil*. 1996;77(6):605-608.
543. Lewis CB, Bottomley JM. *Geriatric Physical Therapy: A Clinical Approach*. Norwalk, CT: Appleton and Lange. 1994:139-186.

544. Licciardone JC, Stoll ST, Cardarelli KM, Gamber RG, Swift JN, Winn WB. A randomized controlled trial of osteopathic manipulative treatment following knee or hip arthroplasty. *J Am Osteopath Assoc*. 2004;104(5):193-202.
545. Lichtenberg PA, Nanna M. The role of cognition in predicting activities of daily living and ambulation functioning in the oldest old rehabilitation patients. *Rehabilitation Psychology*. 1994;39(4) 251-262.
546. Lieberman D, Figer M, Lieberman D. Inpatient rehabilitation outcome after hip fracture surgery in elderly patients: a prospective cohort study of 946 patients. *Arch Phys Med Rehabil*. 2006;87(2):167-171.
547. Lieberman D, Galinsky D, Fried V, et al. Factors affecting the results of the clock drawing test in elderly patients hospitalized for physical rehabilitation. *Int J Geriatr Psychiatry*. 1999;14(5):325-330.
548. Lieberman D, Lieberman D. Rehabilitation after proximal femur fracture surgery in the oldest old. *Arch Phys Med Rehabil*. 2002;83(10):1360-1363.
549. Lieberman D, Lieberman D. Rehabilitation following hip fracture surgery: a comparative study of females and males. *Disabil Rehabil*. 2004;26(2):85-90.
550. Lieberman D, Lieberman D. Rehabilitation following stroke in patients aged 85 and above. *J Rehabil Res Dev*. 2005;42(1):47-53.
551. Lin JH. Influence of admission functional status on functional gain and efficiency of rehabilitation in first time stroke patients. *Kaohsiung J Med Sci*. 2001;17(6):312-318.
552. Lin JH, Chang CM, Liu CK, Huang MH, Lin YT. Efficiency and effectiveness of stroke rehabilitation after first stroke. *J Formos Med Assoc*. 2000;99(6):483-490.
553. Lin JH, Hsiao SF, Chang CM, Huang MH, Liu CK, Lin YT. Factors influencing functional independence outcome in stroke patients after rehabilitation. *Kaohsiung J Med Sci*. 2000;16(7):351-359.
554. Lin JH, Hsiao SF, Liu CK, Lin YT. Rehabilitation fees, length of stay and efficiency for hospitalized stroke patients: a preliminary study based on function-related groups. *Kaohsiung J Med Sci*. 2001;17(9):475-483.
555. Lin JH, Hsieh CL, Hsiao SF, Huang MH. Predicting long-term care institution utilization among post-rehabilitation stroke patients in Taiwan: a medical centre-based study. *Disabil Rehabil*. 2001;23(16):722-730.
556. Lin JH, Hsieh CL, Lo SK, Hsiao SF, Huang MH. Prediction of functional outcomes in stroke inpatients receiving rehabilitation. *J Formos Med Assoc*. 2003;102(10):695-700.
557. Lin JH, Huang MH, Liu CK, Lin YT, Lee CH. The relation between admission balance and functional outcomes following stroke rehabilitation: a medical center based study. *Kaohsiung J Med Sci*. 1999;15(8):491-497.
558. Lin JH, Lo SK, Chang YY, Su CY, Liu CK. Validation of comprehensive assessment of activities of daily living in stroke survivors. *Kaohsiung J Med Sci*. 2004;20(6):287-294.



559. Lin JH, Wei SH, Liu CK, Huang MH, Lin YT. The influence of rehabilitation therapy on the prognosis for stroke patients – a preliminary study. *Kaohsiung J Med Sci*. 1999;15(5):280-291.
560. Lin KH, Wu SC, Hsiung CL, et al. Functional independence of residents in urban and rural long-term care facilities in Taiwan. *Disabil Rehabil*. 2004;26(3):176-181.
561. Linacre JM. Detecting multidimensionality: which residual data-type works best? *Journal of Outcome Measurement*. 1998;2(3):266-83.
562. Linacre JM, Heinemann AW, Wright BD, Granger CV, Hamilton BB. The structure and stability of the Functional Independence Measure. *Archives of Physical Medicine and Rehabilitation*. 1994;75(2):127-132.
563. Linn RT, Blair RS, Granger CV, Harper DW, O'Hara PA, Maciura E. Does the functional assessment measure (FAM) extend the functional independence measure (FIM) instrument? A rasch analysis of stroke inpatients. *J Outcome Meas*. 1999;3(4):339-359.
564. Lippert-Gruner M, Wedekind C, Klug N. Outcome of prolonged coma following severe traumatic brain injury. *Brain Inj*. 2003;17(1):49-54.
565. Liu C, Playford ED, Thompson AJ. Does neurorehabilitation have a role in relapse-remitting multiple sclerosis? *J Neurol*. 2003;250(10):1214-1218.
566. Liu M, Domen K, Chino N. Comorbidity measures for stroke outcome research: a preliminary study. *Archives of Physical Medicine and Rehabilitation*. 1997;78(2):166-172.
567. Liu M, Tsuji T, Higuchi Y, Domen K, Tsujiuchi K, Chino N. Osteoporosis in hemiplegic stroke patients as studied with dual-energy X-ray absorptiometry. *Arch Phys Med Rehabil*. 1999;80(10):1219-1226.
568. Liu M, Tsuji T, Tsujiuchi K, Chino N. Comorbidities in stroke patients as assessed with a newly developed comorbidity scale. *Am J Phys Med Rehabil*. 1999;78(5):416-424.
569. Livingston DH, Lavery RF, Mosenthal AC, et al. Recovery at one year following isolated traumatic brain injury: a Western Trauma Association prospective multicenter trial. *J Trauma*. 2005;59(6):1298-1304; discussion 1304.
570. Long WB, Sacco WJ, Coombes SS, Copes WS, Bullock A, Melville JK. Determining normative standards for Functional Independence Measure transitions in rehabilitation. *Archives of Physical Medicine and Rehabilitation*. 1994;75(2):144-148.
571. Lowthian P, Disler P, Ma S, Eagar K, Green J, de Graaf S. The Australian National Sub-acute and Non-acute Patient Casemix Classification (AN-SNAP): its application and value in a stroke rehabilitation programme. *Clin Rehabil*. 2000;14(5):532-537.
572. Lu AC, Yarkony GM. Benefits of rehabilitation for traumatic spinal cord injury: a case report. *Journal of Spinal Cord Medicine*. 1996; 19(1):17-19.

573. Luk JK, Cheung RT, Ho SL, Li L. Does age predict outcome in stroke rehabilitation? A study of 878 Chinese subjects. *Cerebrovasc Dis*. 2006;21(4):229-234.
574. Lum PS, Burgar CG, Shor PC, Majmundar M, Van der Loos M. Robot-assisted movement training compared with conventional therapy techniques for the rehabilitation of upper-limb motor function after stroke. *Arch Phys Med Rehabil*. 2002;83(7):952-959.
575. Lundgren-Nilsson A, Grimby G, Ring H, et al. Cross-cultural validity of functional independence measure items in stroke: a study using Rasch analysis. *J Rehabil Med*. 2005;37(1):23-31.
576. Lundgren-Nilsson A, Tennant A. Past and present issues in Rasch analysis: the Functional Independence Measure (FIM®) revisited. *J Rehabil Med*. 2011;43:884-891.
577. Lundgren-Nilsson A, Tennant A, Grimby G, Sunnerhagen KS. Cross-diagnostic validity in a generic instrument: an example from the Functional Independence Measure in Scandinavia. *Health Qual Life Outcomes*. 2006;4(1):55.
578. Lutz BJ. Determinants of discharge destination for stroke patients. *Rehabil Nurs*. 2004;29(5):154-163.
579. Lysack CL, MacNeill SE, Lichtenberg PA. The functional performance of elderly urban African-American women who return home to live after medical rehabilitation. *Am J Occup Ther*. 2001;55(4):433-440.
580. Lysack CL, Neufeld S. Occupational therapist home evaluations: inequalities, but doing the best we can? *Am J Occup Ther*. 2003;57(4):369-379.
581. Mackintosh SF, Hill K, Dodd KJ, Goldie P, Culham E. Falls and injury prevention should be part of every stroke rehabilitation plan. *Clin Rehabil*. 2005;19(4):441-451.
582. MacNeill SE, Lichtenberg PA. Home alone: the role of cognition in return to independent living. *Arch Phys Med Rehabil*. 1997;78(7):755-758.
583. Mammi P, Zaccaria B, Franceschini M. Early rehabilitative treatment in patients with traumatic brain injuries: outcome at one-year follow-up. *Eura Medicophys*. 2006;42(1):17-22.
584. Mancuso M, Smith P, Illig S, et al. Satisfaction with medical rehabilitation in patients with orthopedic impairment. *Arch Phys Med Rehabil*. Sep 2003;84(9):1343-1349.
585. Mann WC, Hurren D, Tomita M. Assistive devices used by home-based elderly persons with arthritis. *American Journal of Occupational Therapy*. 1995;49(8):810-820.
586. Mann WC, Hurren D, Tomita M. Comparison of assistive device use and needs of home-based older persons with different impairments. *American Journal of Occupational Therapy*. 1993;47(11):980-987.
587. Mann WC, Hurren D, Tomita M, Charvat B. An analysis of problems with walkers encountered by elderly persons. *Physical and Occupational Therapy in Geriatrics*. 1995;13(1/2):1-23.

588. Mann WC, Hurren D, Tomita M, Charvat BA. A follow-up study of older stroke survivors living at home. *Topics in Geriatric Rehabilitation*. 1995;11(1):52-66.
589. Mann WC, Ottenbacher KJ, Fraas L, Tomita M, Granger CV. Effectiveness of assistive technology and environmental interventions in maintaining independence and reducing home care costs for the frail elderly. A randomized controlled trial. *Arch Fam Med*. 1999;8(3):210-217.
590. Mansson E, Lexell J. Performance of activities of daily living in multiple sclerosis. *Disabil Rehabil*. 2004;26(10):576-585.
591. Marciniak C, Sorosky S, Hynes C. Acute flaccid paralysis associated with West Nile virus: motor and functional improvement in 4 patients. *Arch Phys Med Rehabil*. 2004;85(12):1933-1938.
592. Marciniak CM, Sliwa JA, Spill G, Heinemann AW, Semik PE. Functional outcome following rehabilitation of the cancer patient. *Archives of Physical Medicine and Rehabilitation*. 1996;77(1):54-57.
593. Marino RJ, Crozier KS. Neurological examination and functional assessment after spinal cord injury. *Physical Medicine and Rehabilitation Clinics of North America*. 1992;3(4):829-852.
594. Marino RJ, Graves DE. Metric Properties of the ASIA motor score: subscales improve correlation with functional activities. *Arch Phys Med Rehabil*. 2004;85(11):1804-1810.
595. Marino RJ, Huang M, Knight P, Herbison GJ, Ditunno JF, Segal M. Assessing self-care status in quadriplegia: comparison of the Quadriplegia Index of Function (QIF) and the Functional Independence Measure (FIM). *Paraplegia*. 1993;31(4):225-233.
596. Marino RJ, Rider-Foster D, Maissel G, Ditunno JF. Superiority of motor level over single neurological level of categorizing tetraplegia. *Paraplegia*. 1995;33(9):510-513.
597. Marino RJ, Stineman MG. Functional assessment in spinal cord injury. In: Ditunno JF, ed. *Topics in Spinal Cord Injury Rehabilitation*. Gaithersburg, MD: Aspen Publishers, Inc. 1996;1(4):32-45.
598. Maritime K, Robinson RG. The effect of early versus late antidepressant treatment on physical impairment associated with poststroke depression: is there a time related therapeutic window? *J Nervy Mint Dis*. 2003;191(10):645-652.
599. Marosszeky JE. The Uniform Data System from an international perspective: the view from Australia. *Physical Medicine and Rehabilitation Clinics of North America*. 1993;4(3):571-585.
600. Marshall SC, Heisel B, Grinnell D. Validity of the PULSES profile compared with the Functional Independence Measure for measuring disability in a stroke rehabilitation setting. *Arch Phys Med Rehabil*. 1999;80(7):760-765.
601. Martin MJ, Mulleniz PS, Steele SR, et al. Functional outcome after blunt and penetrating carotid artery injuries: analysis of the National Trauma Data Bank. *J Trauma*. 2005;59(4):860-864.

602. Martin MJ, Weng J, Demetriades D, Salim A. Patterns of injury and functional outcome after hanging: analysis of the National Trauma Data Bank. *Am J Surg*. 2005;190(6):836-840.
603. Masanic CA, Bayley MT, Van Reekum R, Simard M. Open-label study of donepezil in traumatic brain injury. *Arch Phys Med Rehabil*. 2001;82(7):896-901.
604. Mason M, Bell J. Functional outcomes of rehabilitation in the frail elderly: a two-year retrospective review. *Perspectives*. 1994;18(2):7-9.
605. Mauthe RW, Haaf DC, Hayn P, Krall JM. Predicting discharge destination of stroke patients using a mathematical model based on six items from the Functional Independence Measure. *Archives of Physical Medicine and Rehabilitation*. 1996;77:10-13.
606. Mayer M, Bazala J, Muchova B, Stranska H. [Rehabilitation of patients with multiple sclerosis]. *Cas Lek Cesk*. 2001;140(4):118-121.
607. Mayotte JM, Graham R, Mahler L, Young BL, Goodnight S. Impact of nosocomial infection on length of stay and functional improvement among patients admitted to an acute rehabilitation unit. *Infect Control Hosp Epidemiology*. 2001;22(2):83-87.
608. Mazzini L, Campini R, Angelino E, Rognone F, Pastore I, Oliveri G. Posttraumatic hydrocephalus: a clinical, neuroradiologic, and neuropsychologic assessment of long-term outcome. *Arch Phys Med Rehabil*. 2003;84(11):1637-1641.
609. Mazzini L, Cossa FM, Angelino E, Campini R, Pastore I, Monaco F. Posttraumatic epilepsy: neuroradiologic and neuropsychological assessment of long-term outcome. *Epilepsia*. 2003;44(4):569-574.
610. Mazzini L, Zaccala M, Gareri F, Giordano A, Angelino E. Long-latency auditory-evoked potentials in severe traumatic brain injury. *Arch Phys Med Rehabil*. 2001;82(1):57-65.
611. McColl MA, Friedland J. The effects of age and disability on social support. *International Journal of Rehabilitation Research*. 1995;18(4):325-340.
612. McGillis Hall L, Doran D, Baker GR, et al. Nurse staffing models as predictors of patient outcomes. *Med Care*. 2003;41(9):1096-1109.
613. McKinley W, Meade MA, Kirshblum S, Barnard B. Outcomes of early surgical management versus late or no surgical intervention after acute spinal cord injury. *Arch Phys Med Rehabil*. 2004;85(11):1818-1825.
614. McKinley WO, Huang ME, Brunsvold KT. Neoplastic versus traumatic spinal cord injury: an outcome comparison after inpatient rehabilitation. *Arch Phys Med Rehabil*. 1999;80(10):1253-1257.
615. McKinley WO, Kolakowsky SA, Kreutzer JS. Substance abuse, violence, and outcome after traumatic spinal cord injury. *Am J Phys Med Rehabil*. 1999;78(4):306-312.

616. McKinley WO, Tewksbury MA, Mujteba NM. Spinal stenosis vs traumatic spinal cord injury: a rehabilitation outcome comparison. *J Spinal Cord Med.* 2002;25(1):28-32.
617. McNaughton HK, Weatherall M, McPherson KM. Functional measures across neurologic disease states: analysis of factors in common. *Arch Phys Med Rehabil.* 2005;86(11):2184-2188.
618. McPherson KM, Pentland B. Disability in patients following traumatic brain injury – which measure? *Int J Rehabil Res.* 1997;20(1):1-10.
619. McPherson KM, Pentland B, Cudmore SF, Prescott RJ. An inter-rater reliability study of the Functional Assessment Measure (FIM+FAM). *Disabil Rehabil.* 1996;18(7):341-347.
620. Melchiorre PJ, Findley T, Boda W. Functional outcome and comorbidity indexes in the rehabilitation of the traumatic versus the vascular unilateral lower limb amputee. *American Journal of Physical Medicine and Rehabilitation.* 1996;75:9-14.
621. Mellick D, Gerhart KA, Whiteneck GG. Understanding outcomes based on the post-acute hospitalization pathways followed by persons with traumatic brain injury. *Brain Inj.* 2003;17(1):55-71.
622. Menard MR, Hoens AM. Objective evaluation of functional capacity: medical, occupational, and legal settings. *Journal of Orthopaedic and Sports Physical Therapy.* 1994;19(5):249-260.
623. Mendelsohn ME, Leidl DS, Overend TJ, Petrella RJ. Specificity of functional mobility measures in older adults after hip fracture: a pilot study. *Am J Phys Med Rehabil.* 2003;82(10):766-774.
624. Menter RR, Whiteneck GC, Cheliefue SW, et al. Impairment, disability, handicap and medical expenses of persons aging with spinal cord injury. *Paraplegia.* 1991;29(9):613-619.
625. Merbitz C, Morris J, Grip JC. Ordinal scales and foundations of misinference. *Archives of Physical Medicine and Rehabilitation.* 1989;70(4):308-312.
626. Meythaler JM, DeVivo MJ, Braswell WC. Rehabilitation outcomes of patients who have developed Guillain-Barre syndrome. *American Journal of Physical Medicine and Rehabilitation.* 1997;76(5):411-419.
627. Meythaler JM, Hazelwood J, DeVivo MJ, Rosner M. Elevated liver enzymes after nontraumatic intracranial hemorrhages. *Arch Phys Med Rehabil.* 1998;79(7):766-771.
628. Middleton JW, Harvey LA, Batty J, Cameron I, Quirk R, Winstanley J. Five additional mobility and locomotor items to improve responsiveness of the FIM in wheelchair-dependent individuals with spinal cord injury. *Spinal Cord.* 2006;44(8):495-504.
629. Middleton JW, Truman G, Geraghty TJ. Neurological level effect on the discharge functional status of spinal cord injured persons after rehabilitation. *Arch Phys Med Rehabil.* 1998;79(11):1428-1432.

630. Miguens AC, Dias AM, Marques E, Amaral M, Amaral MT, Sousa MR. Functional evaluation of patients with cerebrovascular accident. *Arquivos-de-Fisiatria-e-Doencas-Osteo- Articulares*. 1995;2(4):137-146.
631. Minaire P. The use of the International Classification of Impairments, Disabilities and Handicaps (ICIDH) in Rehabilitation. Strasbourg: Council of Europe, 1989.
632. Mingalla S, Krisciunas A. Occupational therapy for patients with spinal cord injury in early rehabilitation. *Medicina (Kaunas)*. 2005;41(10):852-856.
633. Miyai I, Blau AD, Reding MJ, Volpe BT. Patients with stroke confined to basal ganglia have diminished response to rehabilitation efforts. *Neurology*. 1997;48(1):95-101.
634. Miyai I, Redding MJ. Effects of antidepressants on functional recovery following stroke: a double-blind study. *Journal of Neurological Rehabilitation*. 1998;3(1):55-67.
635. Miyai I, Saito T, Nozaki S, Kang J. A pilot study of the effect of L-threodops on rehabilitation outcome of stroke patients. *Neurorehabil Neural Repair*. 2000;14(2):141-147.
636. Miyai I, Suzuki T, Kii K, Kang J, Kubota K. Wallerian degeneration of the pyramidal tract does not affect stroke rehabilitation outcome. *Neurology*. 1998;51(6):1613-1616.
637. Mizrahi EH, Fleissig Y, Arad M, Adunsky A. Plasma homocysteine level and functional outcome of patients with ischemic stroke. *Arch Phys Med Rehabil*. 2005;86(1):60-63.
638. Mokler PJ, Sandstrom R, Griffin M, Farris L, Jones C. Predicting discharge destination for patients with severe motor stroke: important functional tasks. *Neurorehabil Neural Repair*. 2000;14(3):181-185.
639. Montagnani G, Vagheggini G, Panait Vlad E, Berrighi D, Pantani L, Ambrosino N. Use of the Functional Independence Measure in people for whom weaning from mechanical ventilation is difficult. *Phys Ther*. 2011;91(7):1109-1115.
640. Moore DP, Lee MY, Macciocchi SN. Neurorehabilitation outcome in moyamoya disease. *Arch Phys Med Rehabil*. 1997;78(6):672-675.
641. Moorthi S, Schneider WN, Dombovy ML. Rehabilitation outcomes in encephalitis – a retrospective study 1990 – 1997. *Brain Inj*. 1999;13(2):139-146.
642. Morgan P. The relationship between sitting balance and mobility outcome in stroke. *Australian Journal of Physiotherapy*. 1994;40(2):91-96.
643. Morganti B, Scivoletto G, Ditunno P, Ditunno JF, Molinari M. Walking index for spinal cord injury (WISCI): criterion validation. *Spinal Cord*. 2005;43(1):27-33.
644. Mortifee PR, Busser J, Anton HA. The performance of a limited set of items from the Functional Independence Measure for use in acute trauma care and rehabilitation. *Archives of Physical Medicine and Rehabilitation*. 1996;77:436-439.

645. Mosenthal AC, Livingston DH, Lavery RF, et al. The effect of age on functional outcome in mild traumatic brain injury: 6-month report of prospective multicenter trial. *J Trauma*. 2004;56(5):1042-1048.
646. Moussavi RM, Garza HM, Eisle SG, Rodriguez G, Rintala DH. Serum levels of vitamins A, C, and E in persons with chronic spinal cord injury living in the community. *Arch Phys Med Rehabil*. 2003;84(7):1061-1067.
647. Mouvsas SB, Chang VT, Tunkel RS, Shah VV, Ryan LS, Millis SR. Rehabilitation needs of an inpatient medical oncology unit. *Arch Phys Med Rehabil*. 2003;84(11):1642-1646.
648. Mozzoni MP, Bailey JS. Improving training methods in brain injury rehabilitation. *Journal of Head Trauma Rehabilitation*. 1996;11(1):1-17.
649. Muecke L, Shekar S, Dwyer D, Israel E, Flynn JP. Functional screening of lower-limb amputees: a role in predicting rehabilitation outcome? *Archives of Physical Medicine and Rehabilitation*. 1992;73(9):851-858.
650. Muecke L, Shekar S, Dwyer D, Israel E, Flynn JP. Functional screening of lower-limb amputees: a role in predicting rehabilitation outcome? *Archives of Physical Medicine and Rehabilitation*. 1992;73(9):851-858.
651. Mudge S, Rochester L, Recoran A. The effect of treadmill training on gait, balance and trunk control in a hemiplegic subject: a single system design. *Disabil Rehabil*. 2003;25(17):1000-1007.
652. Mukand J, Larkin L, Barrs K, Lublin P. Midodrine for the management of orthostatic hypotension in patients with spinal cord injury: a case report. *Arch Phys Med Rehabil*. 2001;82(5):694-696.
653. Mukand JA, Cai C, Zielinski A, Danish M, Berman J. The effects of dehydration on rehabilitation outcomes of elderly orthopedic patients. *Arch Phys Med Rehabil*. 2003;84(1):58-61.
654. Mukand JA, Giunti EJ. Tizanidine for the treatment of intention myoclonus: a case series. *Arch Phys Med Rehabil*. 2004;85(7):1125-1127.
655. Mukand JA, Guilmette TJ, Allen DG, et al. Dopaminergic therapy with carbidopa L-dopa for left neglect after stroke: a case series. *Arch Phys Med Rehabil*. 2001;82(9):1279-1282.
656. Mulcahey MJ, Betz RR, Smith BT, Weiss AA. A prospective evaluation of upper extremity tendon transfers in children with cervical spinal cord injury. *J Pediatr Orthop*. 1999;19(3):319-328.
657. Mulcahey MJ, Smith BT, Betz RR. Psychometric rigor of the Grasp and Release Test for measuring functional limitation of persons with tetraplegia: a preliminary analysis. *J Spinal Cord Med*. 2004;27(1):41-46.

658. Mulcahey MJ, Smith BT, Betz RR, Weiss AA. Outcomes of tendon transfer surgery and occupational therapy in a child with tetraplegia secondary to spinal cord injury. *American Journal of Occupational Therapy*. 1995;49(7):607-617.
659. Munin MC, Balu GR, Giuliani MJ, Hegde SK. Neurologic recovery and functional improvement after vecuronium-induced quadriplegia. *American Journal of Physical Medicine & Rehabilitation*. 1995;74(5):375-379.
660. Munin MC, Begley A, Skidmore ER, Lenze EJ. Influence of rehabilitation site on hip fracture recovery in community-dwelling subjects at 6-month follow-up. *Arch Phys Med Rehabil*. 2006;87(7):1004-1006.
661. Munin MC, Rudy TE, Glynn NW, Crossett LS, Rubash HE. Early inpatient rehabilitation after elective hip and knee arthroplasty. *JAMA*. 1998;279(11):847-852.
662. Munin MC, Seligman K, Dew MA, et al. Effect of rehabilitation site on functional recovery after hip fracture. *Arch Phys Med Rehabil*. 2005;86(3):367-372.
663. Murray PK, Dawson NV, Thomas CL, Cebul RD. Are we selecting the right patients for stroke rehabilitation in nursing homes? *Arch Phys Med Rehabil*. 2005;86(5):876-880.
664. Musicco M, Emberi L, Nappi G, Caltagirone C. Early and long-term outcome of rehabilitation in stroke patients: the role of patient characteristics, time of initiation, and duration of interventions. *Arch Phys Med Rehabil*. 2003;84(4):551-558.
665. Muslumanoglu L, Aki S, Ozturk Y, et al. Motor, sensory and functional recovery in patients with spinal cord lesions. *Spinal Cord*. 1997;35(6):386-389.
666. Nagano K. [Appropriate outcome measures for evaluating change in activities of daily living of elderly residents]. *Nippon Koshu Eisei Zasshi*. 2002;49(2):76-87.
667. Nance PW, Huff, FJ, Martinez-Arizala A, Ayyoub Z, Chen D, Bian A, Stamler D. Efficacy and safety study of arbaclofen placarbil in patients with spasticity due to spinal cord injury. *Spinal Cord*. 2011;49(9):974-80.
668. Nanna MJ, Sawilowsky SS. Analysis of Likert Scale data in disability and medical rehabilitation research. *Psychological Methods*. 1998;3(1):55-67.
669. Naps K, Guru A, Civic R, Sara AJ. The relationship between physical impairment and disability during stroke rehabilitation: effect of cognitive status. *Int J Rehabil Res*. 2004;27(3):181-184.
670. Neal LJ. Current functional assessment tools: Barthel Index, the Functional Independence Measure (FIM) and PULSES. *Home Healthcare Nurse*. 1998;16(11):766-72.
671. Nelson DL, Melville LL, Wilkerson JD, Madness RA, Grech JL, Rosenberg JA. Interrater reliability, concurrent validity, responsiveness, and validity of the Melville-Nelson Self-Care Assessment. *Am J Occup Ther*. 2002;56(1):51-59.
672. Nemchausky BA, Ubilluz RM. Upper extremity neuropathies in patients with spinal cord injuries. *Journal of Spinal Cord Medicine*. 1995;18(2):95-97.



673. Neufeld S, Lysack C. Allocation of the rehabilitation services: who gets a home evaluation. *Am J Occup Ther*. 2004;58(6):630-638.
674. New PW. Functional outcomes and disability after nontraumatic spinal cord injury rehabilitation: results from a retrospective study. *Arch Phys Med Rehabil*. 2005;86(2):250-261.
675. Ng YS, Stein J, Salles SS, Black-Schaffer RM. Clinical characteristics and rehabilitation outcomes of patients with posterior cerebral artery stroke. *Arch Phys Med Rehabil*. 2005;86(11):2138-2143.
676. Nguyen-Oghalai TU, Ottenbacher KJ, Granger CV, Goodwin JS. Impact of osteoarthritis on the rehabilitation of patients following a stroke. *Arthritis Rheum*. 2005;53(3):383-387.
677. Nicholas R, Playford ED, Thompson AJ. A retrospective analysis of outcome in severe Guillain-Barre syndrome following combined neurological and rehabilitation management. *Disabil Rehabil*. 2000;22(10):451-455.
678. Nichols DS, Miller L, Colby LA, Pease WS. Sitting balance: its relation to function in individuals with hemiparesis. *Archives of Physical Medicine and Rehabilitation*. 1996;77(9):865-869.
679. Nick TG, Williams JM, Barker JR. Descriptive and graphical strategies for assessing change: a case study of functional status in stroke patients. *Top Health Inf Manage*. 1998;18(3):8-17.
680. Nilsson AL, Sunnerhagen KS, Grimby G. Scoring alternatives for FIM in neurological disorders applying Rasch analysis. *Acta Neurol Scand*. 2005;111(4):264-273.
681. Nilsson L, Carlsson J, Danielsson A, et al. Walking training of patients with hemiparesis at an early stage after stroke: a comparison of walking training on a treadmill with body weight support and walking training on the ground. *Clin Rehabil*. 2001;15(5):515-527.
682. Nir Z, Zolotogorsky Z, Sugarman H. Structured nursing intervention versus routine rehabilitation after stroke. *Am J Phys Med Rehabil*. 2004;83(7):522-529.
683. Nirula R, Brasel K. Do trauma centers improve function outcomes: a national trauma databank analysis? *J Trauma*. 2006;61(2):268-271.
684. Nissen MI. Emphasis placed by Cedar Court Physical Rehabilitation Hospital on the Functional Independence Measure as the overall index for successful rehabilitation. *Australian Clinical Review*. 1989;9(1):36-38.
685. Nolf G, D'Aniello AM, Muschera R, Giaquinto S. The aftermath of rehabilitation for patients with severe stroke. *Acta Neurol Scand*. 2003;107(4):281-284.
686. Nosper M. [Can the classification of patients for phase B, C, and D of neurological rehabilitation be done by FIM™ scores?]. *Rehabilitation (Suttg)*. 2002;41(1):31-39.

687. Novak P, Vidmar G, Kuret Z, Bizovičar N. Rehabilitation of critical illness polyneuropathy and myopathy patients: an observational study. *Int J Rehabil Res.* 2011;34(4):336-42.
688. Nyein K, McMichael L, Turner-Stokes L. Can a Barthel score be derived from the FIM? *Clin Rehabil.* 1999;13(1):56-63.
689. Nyein K, Turner-Stokes L, Robinson I. The Northwick Park Care Needs Assessment (APCNA): a measure of community care needs: sensitivity to change during rehabilitation. *Clin Rehabil.* 1999;13(6):482-491.
690. Oczkowski WJ, Barreca S. The Functional Independence Measure: its use to identify rehabilitation needs in stroke survivors. *Archives of Physical Medicine and Rehabilitation.* 1993;74(12):1291-1294.
691. Oczkowski WJ, Barreca S. Neural network modeling accurately predicts the functional outcome of stroke survivors with moderate disabilities. *Arch Phys Med Rehabil.* 1997;78(4):340-345.
692. O'Dell MW, Barr K, Spanier D, Warnick RE. Functional outcome of inpatient rehabilitation in persons with brain tumors. *Arch Phys Med Rehabil.* 1998;79(12):1530-1534.
693. O'Dell MW, Crawford A, Bohi ES, Bonner FJ Jr. Disability in persons hospitalized with AIDS. *American Journal of Physical Medicine and Rehabilitation.* 1991;70(2):91-95.
694. O'Dell MW, Jasin P, Lyons N, Stivers M, Meszaros F. Standardized assessment instruments for minimally-responsive, brain-injured patients. *NeuroRehabilitation.* 1996;6(1):45-55.
695. O'Dell ME, Watanabe TK, De Roos ST, Kager C. Functional outcome after inpatient rehabilitation in persons with subarachnoid hemorrhage. *Arch Phys Med Rehabil.* 2002;83(5):678-682.
696. Oden KE, Keborkian CG, Levy JK. Rehabilitation of the post-cardiac surgery stroke patient: analysis of cognitive and functional assessment. *Arch Phys Med Rehabil.* 1998;79(1):67-71.
697. Odderson IR, Keaton JC, McKenna BS. Swallow management patients on an acute stroke pathway - quality is cost effective. *Archives of Physical Medicine & Rehabilitation.* 1995; 76(12):1130-1133.
698. O'Donnell JC, Hamilton BB. Stroke rehabilitation management in the Department of Veterans Affairs: impact of patient referral source on outcomes. *Arch Phys Med Rehabil.* 1997;78(9):929-937.
699. Olkkonen S, Lahdenranta U, Slati P, Honkanen R. Bicycle accidents often cause disability - an analysis of medical and social consequences of nonfatal bicycle accidents. *Scandinavian Journal of Social Medicine.* 1993; 21(2):98-106.

700. Ono T, Hori K, Ikebe K, Nokubi T, Nago S, Kookaburra I. Factors influencing eating ability of old in-patients in a rehabilitation hospital in Japan. *Gerodontology*. 2003;20(1):24-31.
701. Ostir GV, Smith PM, Smith D, Rice JL, Ottenbacher KJ. Orthopedic impairment after medical rehabilitation: functional performance and satisfaction with community participation. *Am J Phys Med Rehabil*. 2005;84(10):763-769.
702. Ota T, Akaboshi K, Nagata M, et al. Functional assessment of patients with spinal cord injury: measured by the motor score and the Functional Independence Measure. *Spinal Cord*. 1996;34(9):531-535.
703. O'Toole DM, Goldberg RT, Ryan B. Functional changes in vascular amputee patients: evaluation by Barthel Index, PULSES profile and ESCROW scale. *Archives of Physical Medicine & Rehabilitation*. 1985;66(8):508-511.
704. O'Toole DM, Golden AM. Evaluating cancer patients for rehabilitation potential. *Western Journal of Medicine*. 1991;155(4):384-387.
705. Ottenbacher KJ, Gonzales VA, Smith PM, Illig SB, Fiedler RC, Granger CV. Satisfaction with medical rehabilitation in patients with cerebrovascular impairment. *Am J Phys Med Rehabil*. Dec 2001;80(12):876-884.
706. Ottenbacher KJ, Hsu Y, Granger CV, Fiedler RC. The reliability of the Functional Independence Measure: a quantitative review. *Archives of Physical Medicine and Rehabilitation*. 1996;77(12):1226-1232.
707. Ottenbacher KJ, Mann WC, Granger CV, Tomita M, Hurren D, Charvat B. Inter-rater agreement and stability of functional assessment in the community-based elderly. *Archives of Physical Medicine and Rehabilitation*. 1994;75(12):1297-1301.
708. Ottenbacher KJ, Msall ME, Lyon NR, Duffy LC, Granger CV, Braun S. Interrater agreement and stability of the Functional Independence Measure for Children (WeeFIM): use in children with developmental disabilities. *Arch Phys Med Rehabil*. 1997;78(12):1309-1315.
709. Ottenbacher KJ, Smith PM, Illig SB, Fiedler RC, Gonzales V, Granger CV. Characteristics of persons rehospitalized after stroke rehabilitation. *Arch Phys Med Rehabil*. Oct 2001;82(10):1367-74.
710. Ottenbacher KJ, Taylor ET, Msall ME, et al. The stability and equivalence reliability of the functional independence measure for children (WeeFIM). *Dev Med Child Neurol*. 1996;38(10):907-916.
711. Ouslander JG, Griffiths PC, McConnell E, Riolo L, Kutner M, Schnelle J. Functional incidental training: a randomized, controlled, crossover trial in Veterans Affairs nursing homes. *J Am Geriatr Soc*. 2005;53(7):1091-1100.
712. Owen DC, Getz PA, Bulla S. A comparison of characteristics of patients with completed stroke: those who achieve continence and those who do not. *Rehabilitation Nursing*. 1995;20(4):197-203.

713. Ozdemir F, Birtane M, Tabatabaei R, Kokino S, Ekuklu G. Comparing stroke rehabilitation outcomes between acute inpatient and nonintense home settings. *Arch Phys Med Rehabil.* 2001;82(10):1375-1379.
714. Ozgocmen S, Bulut S, Ilhan N, Gulkesen A, Ardicoglu O, Ozkan Y. Vitamin D deficiency and reduced bone mineral density in multiple sclerosis: effect of ambulatory status and functional capacity. *J Bone Miner Metab.* 2005;23(4):309-313.
715. Padua L, Rendeli C, Rabini A, Girardi E, Tonali P, Salvaggio E. Health-related quality of life and disability in young patients with spina bifida. *Arch Phys Med Rehabil.* 2002;83(10):1384-1388.
716. Panesar BS, Morrison P, Hunter J. A comparison of three measures of progress in early lower limb amputee rehabilitation. *Clin Rehabil.* 2001;15(2):157-171.
717. Paolinelli C, Gonzalez P, Doniez ME, Donosos T, Salinas V. [Instrument for the functional evaluation of disability rehabilitation. Study on the reliability and clinical experience with the use of functional independence measure]. *Rev Med Chil.* 2001;129(1):23-31.
718. Parfenchuck TA, Parziale JR, Liberman JR, Butcher RP, Ahern DK. The evolution of an acute care hospital unit to a DRG-exempt rehabilitation unit. A preliminary communication. *American Journal of Physical Medicine and Rehabilitation.* 1990;69(1):11-15.
719. Parsch D, Mikut R, Abel R. Postacute management of patients with spinal cord injury due to metastatic tumour disease: survival and efficiency of rehabilitation. *Spinal Cord.* 2003;41(4):205-210.
720. Passler MA, Riggs RV. Positive outcomes in traumatic brain injury-vegetative state: patients treated with bromocriptine. *Arch Phys Med Rehabil.* 2001;82(3):311-315.
721. Patti F, Ciancio MR, Cacopardo M, et al. Effects of a short outpatient rehabilitation treatment on disability of multiple sclerosis patients – a randomized controlled trial. *J Neurol.* 2003;250(7):861-866.
722. Peel C, Utsey C. Role of endurance exercise in older persons with physical disabilities resulting from CVA: a case report. *Issues on Aging.* 18(1):19-24.
723. Pentland W, McColl MA, Rosenthal C. The effect of aging and duration of disability on long term health outcomes following spinal cord injury. *Paraplegia.* 1995; 33(7):367-373.
724. Perino C, Rago R, Cicolini A, Torta R, Monaco F. Mood and behavioural disorders following traumatic brain injury: clinical evaluation and pharmacological management. *Brain Inj.* 2001;15(2):139-148.
725. Peron EP, Gray SL, Hanlon JT. Medication use and functional status decline in older adults: a narrative review. *Am J Geriatr Pharmacother.* 2011;9(6):378-91.
726. Perron AD, Sing RF, Branas CC, Huynh T. Predicting survival in pediatric trauma patients receiving cardiopulmonary resuscitation in the prehospital setting. *Prehosp Emerg Care.* 2001;5(1):6-9.

727. Perry J. Rehabilitation of the neurologically disabled patient: principles, practice, and scientific basis. *Journal of Neurosurgery*. 1983;58(6):799-816.
728. Perry SB, Marchetti GF, Wagner S, Wilton W. Predicting caregiver assistance required for sit-to-stand following rehabilitation for acute stroke. *J Neurol Phys Ther*. 2006;30(1):2-11.
729. Peter A. Evaluation of the Uniform Data System for Medical Rehabilitation (including the Functional Independence Measure) by Glenrothes Geriatric Day Hospital team. *British Journal of Occupational Therapy*. 1994;57(3):91-94.
730. Petracca G, Teson A, Chemerinski E, Leiguarda R, Starkstein SE. A double-blind placebo-controlled study of clomipramine in depressed patients with Alzheimer's disease. *J Neuropsychiatry Clin Neurosci*. 1996;8(3):270-275.
731. Petrella RJ, Jones TJ. Do patients receive recommended treatment of osteoporosis following hip fracture in primary care? *BMC Fam Pract*. 2006;7:31.
732. Petrella RJ, Overend T, Chesworth B. FIM after hip fracture: is telephone administration valid and sensitive to change? *Am J Phys Med Rehabil*. 2002;81(9):639-644.
733. Petrilli S, Durufle A, Nicholas B, Pinel JF, Kerdoncuff V, Gallien P. [Hemiplegia and return to domicile]. *Ann Readapt Med Phys*. 2002;45(2):69-76.
734. Petruseviciene D, Krisciunas A. [Influence of factors on independence of patients after stroke in early rehabilitation stage]. *Medicina (Kaunas)*. 2005;41(8):655-660.
735. Petruseviciene D, Krisciunas A. [Influence of factors on occupational therapy efficiency for patients after stroke]. *Medicina (Kaunas)*. 2005;41(4):320-324.
736. Peurala SH, Tarkka IM, Pitkanen K, Sivenius J. The effectiveness of body weight-supported gait training and floor walking in patients with chronic stroke. *Arch Phys Med Rehabil*. 2005;86(8):1557-1564.
737. Phipps MA, Kelly-Hayes M. Rehabilitation of older adults. In: Baines EM. ed. *Perspectives on Gerontological Nursing*. Newbury Park: Sage Publications. 1991:357-373.
738. Pinholt EM, Kroenke K, Hanley JF, Kussman MJ, Twyman PL, Carpenter JL. Functional assessment of the elderly. A comparison of standard instruments with clinical judgment. *Archives of Internal Medicine*. 1987;147(3):484-488.
739. Ploughman M, Corbett D. Can forced-use therapy be clinically applied after stroke? An exploratory randomized controlled trial. *Arch Phys Med Rehabil*. 2004;85(9):1417-1423.
740. Priebe MM, Rintala DH. Functional assessment in a community setting. *Topics in Stroke Rehabilitation*. 1994;1(3):16-29.

741. Poduri KR, Cushman LA, Gibson CJ. Inpatient rehabilitation: the correlation between functional gains and appropriateness of admissions. *Int J Rehabil Res.* 1996;19(4):327-332.
742. Pollak N, Rheault W, Stoecker JL. Reliability and validity of the FIM for persons aged 80 years and above from a multilevel continuing care retirement community. *Archives of Physical Medicine and Rehabilitation.* 1996;77:1056-1061.
743. Poon WS, Zhu XL, Ng SC, Wong GK. Predicting one year clinical outcome in traumatic brain injury (TBI) at the beginning of rehabilitation. *Acta Neurochir Suppl.* 2005;93:207-208.
744. Popovic MR, Thrasher TA, Adams ME, Takes V, Zivanovic V, Tonack MI. Functional electrical therapy: retraining grasping in spinal cord injury. *Spinal Cord.* 2006;44(3):143-151.
745. Post MW, Dallmeijer AJ, Angenot EL, van Asbeck FW, van der Woude LH. Duration and functional outcome of spinal cord injury rehabilitation in the Netherlands. *J Rehabil Res Dev.* 2005;42(3 Suppl 1): 75-85.
746. Powell J, Heslin J, Greenwood R. Community based rehabilitation after severe traumatic brain injury: a randomized controlled trial. *J Neurol Neurosurg Psychiatry.* 2002;72(2):193-202.
747. Prasad R, Hellawell DJ, Pentland B. Usefulness of the Functional Independence Measure (FIM), its subscales and individual items as outcome measures in Guillain-Barre Syndrome. *Int J Rehabil Res.* 2001;24(1):59-64.
748. Przbylski BR, Dumont D, Watkins ME, Warren SA, Beaulne AP, Lier DA. Outcomes of enhanced physical and occupational therapy service in a nursing home setting. *Archives of Physical Medicine and Rehabilitation.* 1996;77(6):554-561.
749. Putzke JD, Barrett JJ, Richards JS, DeVivo MJ. Age and spinal cord injury: an emphasis on outcomes among the elderly. *J Spinal Cord Med.* 2003;26(1):37-44.
750. Putzke JD, Hicken BL, Richard JS. Race: predictor versus proxy variable? Outcomes after spinal cord injury. *Arch Phys Med Rehabil.* 2002;83(11):1603-1611.
751. Qiang W, Sonoda S, Suzuki M, Okamoto S, Saitoh E. Reliability and validity of a wheelchair collision test for screening behavioral assessment of unilateral neglect after stroke. *Am J Phys Med Rehabil.* 2005;84(3):161-166.
752. Rabadi MH, Blau A. Admission ambulation velocity predicts length of stay and discharge disposition following stroke in an acute rehabilitation hospital. *Neurorehabil Neural Repair.* 2005;19(1):20-26.
753. Rabadi MH, Rabadi FM. Comparison of the action research arm test and the Fugl-Meyer assessment as measures of upper-extremity motor weakness after stroke. *Arch Phys Med Rehabil.* 2006;87(7):962-966.
754. Rabinowitz JC, Craik FIM. Prior retrieval effects in young and old adults. *Journal of Gerontology.* 1986;41(3):368-75.

755. Rao N, Cahr D, Gnatz S. Rehabilitation outcomes of 5 patients with sever West Nile virus infection: a case series. *Arch Phys Med Rehabil.* 2005;86(3):449-452.
756. Ravaud JF, Delcey M, Yelnik A. Construct validity of the functional independence measure (FIM): questioning the unidimensionality of the scale and the "value" of FIM scores. *Scand J Rehabil Med.* 1999;31(1):31-41.
757. Rawl SM, Easton KL, Kwiatkowski S, Zemen D, Burczyk B. Effectiveness of a nurse-managed follow-up program for rehabilitation patients after discharge. *Rehabil Nurs.* 1998;23(4):204-209.
758. Reiskind M, Kirshblum SC, Maney M, Johnson MV. The Functional Independence Measure as a predictor of falls in an inpatient stroke rehabilitation. *Archives of Physical Medicine and Rehabilitation.* 1992;73(10):965. (Abstract)
759. Rendeli C, Ausili E, Tabacco F, et al. Assessment of health status in children with spina bifida. *Spinal Cord.* 2005;43(4):230-235.
760. Ricci NA, Kubota MT, Cordeiro RC. [Agreement between observations on the functional capacity of home care elderly patients]. *Rev Saude Publica.* 2005;39(4):655-662.
761. Richard R, Eigler M. The Functional Independence Measure (FIM) and functional outcomes after a new approach to axillary burns. *J Burn Care Rehabil.* 2005;26(6):529-531.
762. Richards LG, Latham NK, Jette DU, Rosenberg L, Smout RJ, DeJong G. Characterizing occupational therapy practice in stroke rehabilitation. *Arch Phys Med Rehabil.* 2005;86(12 Suppl 2):S51-S60.
763. Ring H, Bar L, Abboud S. Functional correlates with left-right asymmetry of visual evoked potentials in stroke patients: modeling and experimental results. *Arch Phys Med Rehabil.* 1999;80(2):166-172.
764. Ring H, Feder M, Schwartz J, Samuels G. Functional measures of first-stroke rehabilitation inpatients: usefulness of the Functional Independence Measure total score with a clinical rationale. *Arch Phys Med Rehabil.* 1997;78(6):630-635.
765. Rintala DH, Young ME, Hart KA, Fuhrer MJ. The relationship between the extent of reciprocity with social supporters and measures of depressive symptomatology, impairment, disability, and handicap in persons with spinal cord injury. *Rehabilitation Psychology.* 1994; 39(1):15-27.
766. Rissanen P, Aro S, Sintonen H, Asikainen K, Slati P, Paavolainen P. Costs and cost-effectiveness in hip and knee replacements. A prospective study. *Int J Technol Assess Health Care.* 1997;13(4):575-588.
767. Rockwood K, Stolee P, Fox RA. Use of goal attainment scaling in measuring clinically important change in the frail elderly. *Journal of Clinical Epidemiology.* 1993;46(10):1113-1118.

768. Rolland Y, Pillard F, Lauwers-Cances V, Busquere F, Velals B, Lafont C. Rehabilitation outcome of elderly patients with hip fracture and cognitive impairment. *Disabil Rehabil.* 2004;26(7):425-431.
769. Romberg A, Virtanen A, Ruutiainen J. Long-term exercise improves functional impairment but not quality of life in multiple sclerosis. *J Neurol.* 2005;252(7):839-845.
770. Roques CF, Felez A, Marque P, Chatain M, Condouret J, Tuffery R. [A motor scale for the hemiplegic patients' motor function assessment]. *Annales de Readaptation et de Médecine Physique.* 1997;40(3):147-158.
771. Roth E, Davidoff G, Heughton J, Ardner M. Functional assessment in spinal cord injury: a comparison of the Modified Barthel Index and the 'adapted' Functional Independence Measure. *Clinical Rehabilitation.* 1990;4:277-285.
772. Roth E, Heinemann AW, Lovell LL, Harvey RL, McGuire JR, Diaz S. Impairment and disability: their relation during stroke rehabilitation. *Archives of Physical Medicine and Rehabilitation.* 1998;79(3):329-35.
773. Roth EJ, Lovell L, Harvey RL, Bode RK, Heinemann AW. Stroke rehabilitation: indwelling urinary catheters, enteral feeding tubes, and tracheostomies are associated with resource use and functional outcomes. *Stroke.* 2002;33(7):1845-1850.
774. Rousseaux M, Kozlowski O, Forger J. Efficacy of botulinum toxin A in upper limb function of hemiplegic patients. *J Neurol.* 2002;249(1):76-84.
775. Routhier F, Desrosiers J, Vincent C, Nadeau S. Reliability and validity studies of an obstacle course assessment of wheelchair user performance. *Int J Rehabil Res.* 2005;28(1):49-56.
776. Ruchinskas RA, Singer HK, Repetz NK. Cognitive status and ambulation in geriatric rehabilitation: walking without thinking? *Arch Phys Med Rehabil.* 2000;81(9):1224-1228.
777. Ruchinskas RA, Singer HK, Repetz NK. Clock drawing, clock copying, and physical abilities in geriatric rehabilitation. *Arch Phys Med Rehabil.* 2001;82(7):920-924.
778. Rupright J, Singh A, Stonnington H, et al. Correlation of functional independence measure (FIM) and SPECT lofetamine (I-123) as a predictor of functional return in stroke. *Brain Inj.* 1997;11(1):49-57.
779. Saboe LA, Darrah JM, Pain KS, Guthrie J. Early predictors of functional independence 2 years after spinal cord injury. *Arch Phys Med Rehabil.* 1997;78(6):644-650.
780. Saciri BM, Kos N. Aneurysmal subarachnoid haemorrhage: outcomes of early rehabilitation after surgical repair of ruptured intracranial aneurysms. *J Neurol Neurosurg Psychiatry.* 2002;72(3):334-337.
781. Saikkonen J, Karppi P, Hussko TM, Dahlberg A, Makinen J, Uutela T. Life situation of spinal cord-injured persons in Central Finland. *Spinal Cord.* 2004;42(8):459-465.



782. Sager MA, Dunham NC, Schwantes A, Mecum L, et al. Measurement of activities of daily living in hospitalized elderly: A comparison of self-report and performance-based methods. *Journal of the American Geriatrics Society*. 1992;40(5):457-462.
783. San Segundo RM, Aguilar JJ, Santos F, Usabiaga T. A model for predicting delay in discharge of stroke patients. *Ann Readapt Med Phys*. 2007;50(1):14-9.
784. Sander AM, Fuchs KL, High WM, Jr., Hall KM, Kreutzer JS, Rosenthal M. The Community Integration Questionnaire revisited: an assessment of factor structure and validity. *Arch Phys Med Rehabil*. 1999;80(10):1303-1308.
785. Sandstrom R, Mokler PJ, Hoppe KM. Discharge destination and motor function outcome in severe stroke as measured by the functional independence measure/function-related group classification system. *Arch Phys Med Rehabil*. 1998;79(7):762-765.
786. Sangha H, Lipson D, Foley N, et al. A comparison of the Barthel Index and the Functional Independence Measure as outcome measures in stroke rehabilitation: patterns of disability scale usage in clinical trials. *Int J Rehabil Res*. 2005;28(2):135-139.
787. Sansone GR, Alba A, Frengley JD. Analysis of FIM instrument scores for patients admitted to an inpatient cardiac rehabilitation program. *Arch Phys Med Rehabil*. 2002;83(4):506-512.
788. Sato D, Kaneko F, Okamura H. Reliability and validity of the Japanese-language version of the physical performance test (PPT) battery in chronic pain patients. *Disabil Rehabil*. 2006;28(6):397-405.
789. Sato W, Yanagi H, Yamada K, Suzuki M, Tomura S, Tsuchiya S. [A study on supporting methods for house-bound home care elderly to get independent life (1<sup>st</sup> report) – relation between ADL and daily hours out of bed]. *Nihon Kango Kagakkaishi*. 1997;17(1):66-74.
790. Saverino A, Benevolo E, Ottonello M, Zsirai E, Sessarego P. Falls in a rehabilitation setting: functional independence and fall risk. *Eura Medicophys*. 2006;42(3):179-84.
791. Savic G, Short DJ, Weitzenkamp D, Charlifue S, Gardner BP. Hospital readmissions in people with chronic spinal cord injury. *Spinal Cord*. 2000;38(6):371-377.
792. Schaufele MK, Jacobi E. [Developing a rehabilitation databank for Germany – conceptual considerations exemplified by UDSMR<sub>SM</sub> (Uniform Data System of Medical Rehabilitation)]. *Rehabilitation (Suttg)*. 1999;38(3):149-153.
793. Schemm RL, Gitlin LN. How occupational therapists teach older patients to use bathing and dressing devices in rehabilitation. *American Journal of Occupational Therapy*. 1998;52(4):276-282.
794. Schepers VP, Ketelaar M, Visser-Meily JM, Dekker J, Lindeman E. Responsiveness of functional health status measures frequently used in stroke research. *Disabil Rehabil*. 2006;28(17):1035-1040.

795. Schindl MR, Forstner C, Kern H, Zipko HT, Rupp M, Zifko UA. Evaluation of a German version of the Rivermead Mobility Index (RMI) in acute and chronic stroke patients. *Eur J Neurol*. 2000;7(5):523-528.
796. Schleenbaker RE, McDowell SM, Moore RW, Costich JF, Prater G. Restraint use in inpatient rehabilitation: incidence, predictors, and implications. *Archives of Physical Medicine and Rehabilitation*. 1994;75(4):427-430.
797. Schneider J, Gerrard P, Goldstein R, et al. The impact of comorbidities and complications on burn injury inpatient rehabilitation outcomes. *PM&R*. 2012;5(2):114-121.
798. Schneider JC, Gerrard P, Goldstein R, et al. Predictors of transfer from rehabilitation to acute care in burns. *J Trauma Acute Care Surg*. 2012;73(6):1594-1599.
799. Schneider JC, Tan WH, Goldstein R, et al. Time to rehabilitation in the burn population: Incidence of zero onset days in the UDSMR national dataset. *J Burn Care Res*. 2013;Mar 8 epub ahead of print.
800. Scialla SJ, Cole RP, Bednarz L. Redefining cancer-related asthenia-fatigue syndrome. *J Palliat Med*. 2006;9(4):866-872.
801. Sebastia E, Duarte E, Boza R, et al. Cross-validation of a model for predicting functional status and length of stay in patients with stroke. *J Rehabil Med*. 2006;38(3):204-206.
802. Seel RT, Huang ME, Cifu DX, Kolakowsky-Hayner SA, McKinley WO. Age-related differences in length of stays, hospitalization costs, and outcomes for an injury-matched sample of adults with paraplegia. *J Spinal Cord Med*. 2001;24(4):241-250.
803. Segal ME, Ditunno JF, Staas WE. Inter-institutional agreement of individual Functional Independence Measure (FIM) items measured at two sites on one sample of SCI patients. *Paraplegia*. 1993;31(10):622-631.
804. Segal ME, Gillard M, Schall RR. Telephone and in-person proxy agreement between stroke patients and caregivers for the Functional Independence Measure. *American Journal of Physical Medicine and Rehabilitation*. 1996;75(3):208-212.
805. Segal ME, Schall RR. Determining functional/health status and its relation to disability in stroke survivors. *Stroke*. 1994;25(12):2391-2397.
806. Segal ME, Schall RR. Assessing handicap of stroke survivors. A validation study of the Craig Handicap Assessment and Reporting Technique. *American Journal of Physical Medicine and Rehabilitation*. 1995;74(4):276-286.
807. Seidel GK, Millis SR, Lichtenberg PA, Dijkers M. Predicting bowel and bladder continence from cognitive status in geriatric rehabilitation patients. *Archives of Physical Medicine and Rehabilitation*. 1994;75(5):590-593.
808. Seliger GM, Lichtman SW, Hornstein A. Naltrexone improves several posttraumatic abulia. *Journal of Neurological Rehabilitation*. 1998;12(1):29-31.

809. Seki M, Takenaka A, Nakazawa M, Takahashi H, Chino N. [Examination of living environment upon return to home for patients with cervical spinal cord injury – report of a case]. *Gan To Kagaku Ryoho*. 2002;29 (Suppl 3):522-525.
810. Semlyen JK, Summers SJ, Barnes MP. Traumatic brain injury: efficacy of multidisciplinary rehabilitation. *Arch Phys Med Rehabil*. 1998;79(6):678-683.
811. Seymour DG, Ball AE, Russell EM, Primrose WR, Garratt AM, Crawford JR. Problems in using health survey questionnaires in older patients with physical disabilities. The reliability and validity of the SF-36 and the effect of cognitive impairment. *J Eval Clin Pract*. 2001;7(4):411-418.
812. Shah S, Cooper B. Issues in the choice of activities of daily living assessment. *The Australian Occupational Therapy Journal*. 1993;40(2):77-82.
813. Sharrack B, Hughes RA, Soudain S, Dunn G. The psychometric properties of clinical rating scales used in multiple sclerosis. *Brain*. 1999;122(Pt 1):141-159.
814. Shaughnessy M, Michael KM, Sorkin JD, Macko RF. Steps after stroke: capturing ambulatory recovery. *Stroke*. 2005;36(6):1305-1307.
815. Shectman O, Mann WC, Justiss MD, Tomita M. Grip strength in the frail elderly. *Am J Phys Med Rehabil*. 2004;83(11):819-826.
816. Shelton FD, Volpe BT, Reding M. Motor impairments as a predictor of functional recovery and guide to rehabilitation treatment after stroke. *Neurorehabil Neural Repair*. 2001;15(3):229-237.
817. Shen Q, Cordato D, Chan DK, Hung WT, Karr M. Identifying the determinants of 1-year post-stroke outcomes in elderly patients. *Act Neurol Scand*. 2006;113(2):114-120.
818. Sheng AZ, Shen Q, Cordato D, Zhang YY, Yin Chan DK. Delirium within three days of stroke in a cohort of elderly patients. *J Am Geriatr Soc*. 2006;54(8):1192-1198.
819. Sherer M, Hat T, Nick TG, Whyte J, Thompson RN, Yablon SA. Early impaired self-awareness after traumatic brain injury. *Arch Phys Med Rehabil*. 2003;84(2):168-176.
820. Shiflett SC, Nayak S, Bid C, Miles P, Agostinelli S. Effect of Reiki treatments on functional recovery in patients poststroke rehabilitation: a pilot study. *J Altern Complement Med*. 2002;8(6):755-763.
821. Shiri-Sharvit O, Arad M, Mizrahi EH, Fleissig Y, Adunsky A. The association between psychotropic medication use and functional outcome of elderly hip-fracture patients undergoing rehabilitation. *Arch Phys Med Rehabil*. 2005;86(7):1389-1393.
822. Shoemaker MJ, Mullins-Macritchie M, Bennett J, Vryhof K, Boettger I. Predicting response to rehabilitation in elderly patients with stroke using the orpington prognostic scale and selected clinical variables. *J Geriatr Phys Ther*. 2006;29(2):69-73.
823. Shone SM, Ryan S, Rigby PJ, Jutai JW. Toward a comprehensive evaluation of the impact of electronic aids to daily living: evaluation of consumer satisfaction. *Disabil Rehabil*. 2002;24(1-3):115-125.

824. Siebens H, Andres PL, Pengsheng N, Coster WJ, Haley SM. Measuring physical function in patients with complex medical and postsurgical conditions: a computer adaptive approach. *Am J Phys Med Rehabil.* 2005;84(10):741-748.
825. Silverstein B, Fisher WP, Kilgore KM, Harley JP, Harvey RF. Applying psychometric criteria to functional assessment in medical rehabilitation: II. defining interval measures. *Archives of Physical Medicine and Rehabilitation.* 1992;73:507-518.
826. Simondson JA, Goldie P, Greenwood KM. The Mobility Scale for Acute Stroke Patients: concurrent validity. *Clin Rehabil.* 2003;17(5):558-564.
827. Singh A, Black SE, Herrmann N, et al. Functional and neuroanatomic correlations in poststroke depression: the Sunnybrook Stroke Study. *Stroke.* 2000;31(3):637-644.
828. Singh R, Hunter J, Philip A, Todd I. Predicting those who will walk after rehabilitation in a specialist stroke unit. *Clin Rehabil.* 2006;20(2):149-152.
829. Sipski ML, Jackson AB, Gomez-Marin O, Estores I, Stein A. Effects of gender on neurologic and functional recovery after spinal cord injury. *Arch Phys Med Rehabil.* 2004;85(11):1826-1936.
830. Siu AL, Boockvar KS, Penrod JD, et al. Effect of inpatient quality of care on functional outcomes in patients with hip fracture. *Med Care.* 2006;44(9):862-869.
831. Skinner A, Turner-Stokes L. The use of standardized outcome measures in rehabilitation centres in the UK. *Clin Rehabil.* 2006;20(7):609-615.
832. Silwa JA, Heinemann A, Semik P. Inpatient rehabilitation following brain injury: patient demographics and functional outcomes. *Arch Phys Med Rehabil.* 2005;86(10):1920-1923.
833. Smith BT, Mulcahey MJ, Betz RR. An implantable upper extremity neuroprosthesis in a growing child with C5 spinal cord injury. *Spinal Cord.* 2001;39(2):118-123.
834. Smith PM, Bennett S, Fielder RC, Hamilton BB, Ottenbacher KJ. Intermodal agreement of follow-up telephone functional assessment using the Functional Independence Measure in patients with stroke. *Archives of Physical Medicine and Rehabilitation.* 1996;77(5):431-435.
835. Smith PM, Illig SB, Fielder RC, Hamilton BB, Ottenbacher KJ. Intermodal agreement of follow-up telephone functional assessment using the Functional Independence Measure in patients with stroke. *Arch Phys Med Rehabil.* 1996;77(5):431-435.
836. Smith PM, Ottenbacher KJ, Cranley M, Dittmar SS, Illig SB, Granger CV. Predicting follow-up living setting in patients with stroke. *Arch Phys Med Rehabil.* Jun 2002;83(6):764-770.
837. Smith R. Validation and reliability of the Elderly Mobility Scale. *Physiotherapy.* 1994;80(11):744-747.

838. Smith-Knapp K, Corrigan JD, Arnett JA. Predicting functional independence from neuropsychological tests following traumatic brain injury. *Brain Inj.* 1996;10(9):651-661.
839. Solari A, Filippini G, Gasco P, et al. Physical rehabilitation has a positive effect on disability in multiple sclerosis patients. *Neurology.* 1999;52(1):57-62.
840. Sonoda S, Chino N, Domen K, Saitoh E. Changes in impairment and disability from the third to the sixth month after stroke and its relationship evaluated by an artificial neural network. *Am J Phys Med Rehabil.* 1997;76(5):395-400.
841. Sonoda S, Saitoh E, Nagai S, Kawakita M, Kanada Y. Full-time integrated treatment program, a new system for stroke rehabilitation in Japan: comparison with conventional rehabilitation. *Am J Phys Med Rehabil.* 2004;83(2):88-93.
842. Sonn U, Svensson E. FIM measurement properties and Rasch model details. *Scandinavian Journal of Rehabilitation Medicine.* 1997;29:267-272.
843. Sorbo A, Rydenhag B, Sunnerhagen KS, Blomqvist M, Svensson S, Emanuelson I. Outcome after severe brain damage, what makes the difference? *Brain Inj.* 2005;19(7):493-503.
844. Speed J. Behavioral management of conversion disorder - retrospective study. *Archives of Physical Medicine and Rehabilitation.* 1996;77(2):147-54.
845. Spires MC, Bowden ML, Ahrns KS, Wahl WL. Impact of an inpatient rehabilitation facility on functional outcome and length of stay of burn survivors. *J Burn Care Rehabil.* 2005;26(6):532-538.
846. Spooren AJ, Janssen-Potten YJ, Post MW, Kerckhofs E, Nene A, Seele HA. Measuring change in arm hand skilled performance in persons with a cervical spinal cord injury: responsiveness of the Van Lieshout Test. *Spinal Cord.* 2006;44(12):772-779.
847. Stalp M, Koch C, Regel G, Hretteck C, Pape HE. [Development of a standardized instrument for quantitative and reproducible rehabilitation data assessment after polytrauma (HASPOC)]. *Chirurg.* 2001;72(3):312-318.
848. Stanley RK, Stafford DJ, Rasch E, Rodgers MM. Development of a functional assessment measure for manual wheelchair users. *J Rehabil Res Dev.* 2003;40(4):301-307.
849. Steube D, Wietholter S, Correll C. Prognostic Value of lower limb motor evoked potentials for motor impairment and disability after 8 weeks of stroke rehabilitation – a prospective investigation of 100 patients. *Electromyogr Clin Neurophysiol.* 2001;41(8):463-469.
850. Stevenson VL, Playford ED, Langdon DW, Thompson AJ. Rehabilitation of incomplete spinal cord pathology: factors affecting prognosis and outcome. *J Neurol.* 1996;243(9):644-647.
851. Stilwell P, Stilwell J, Hawley C, Davies C. Measuring outcome in community-based rehabilitation services for people who have suffered traumatic brain injury: the Community Outcome Scale. *Clin Rehabil.* 1998;12(6):521-531.

852. Stineman MG. Function-based classification for stroke rehabilitation and issues of reimbursement: using patient classification systems to scale payment to patient complexity. *Topics in Stroke Rehabilitation*. 1994;1(3):40-50.
853. Stineman MG. Measuring casemix, severity, and complexity in geriatric patients undergoing rehabilitation. *Med Care*. 1997;35(6 Suppl):JS90-105; discussion JS106-102.
854. Stineman MG. The story of function-related groups – please, first do no harm. *Arch Phys Med Rehabil*. 2001;82(4):553-557.
855. Stineman MG. Prospective payment, prospective challenge. *Arch Phys Med Rehabil*. 2002;83(12):1802-1805.
856. Stineman MG, Escarce JJ, Goin JE, Hamilton BB, Granger CV, Williams SV. A case-mix classification system for medical rehabilitation. *Medical Care*. 1994;32(4):366-379.
857. Stineman MG, Escarce JJ, Tassoni CJ, Goin JE, Granger CV, Williams SV. Diagnostic coding and medical rehabilitation length of stay: their relationship. *Arch Phys Med Rehabil*. 1998;79(3):241-248.
858. Stineman MG, Fielder RC, Granger CV, Maislin G. Functional task benchmarks for stroke rehabilitation. *Arch Phys Med Rehabil*. 1998;79(5):497-504.
859. Stineman MG, Goin JE, Granger CV, Fielder R, Williams SV. Discharge motor FIM-function related groups. *Arch Phys Med Rehabil*. 1997;78(9):980-985.
860. Stineman MG, Goin JE, Hamilton BB, Granger CV. Efficiency pattern analysis for medical rehabilitation. *American Journal of Medical Quality*. 1995;10(4):190-203.
861. Stineman MG, Goin JE, Tassoni CJ, Granger CV, Williams SV. Classifying rehabilitation patients by expected functional gain. *Medical Care*. 1997;35(9):963-73.
862. Stineman MG, Granger CV. Epidemiology of stroke-related disability and rehabilitation outcome. *Physical Medicine and Rehabilitation Clinics of North America*. 1991;2:457-471.
863. Stineman MG, Granger CV. Outcome studies and analyses: principles of rehabilitation that influence outcome analyses. In: Felsenthal G, Garrison SJ, Steinberg FU, eds. *Rehabilitation of the Aging and Elderly*. Baltimore: Williams and Wilkins, 1994:511-522.
864. Stineman MG, Granger CV. A modular case-mix classification system for medical rehabilitation illustrated. *Health Care Financ Rev*. 1997;19(1):87-103.
865. Stineman MG, Granger CV. Outcome, efficiency, and time-trend pattern analysis for stroke rehabilitation. *American Journal of Physical Medicine and Rehabilitation*. 1998;77(3):193-201.
866. Stineman MG, Granger CV, Hamilton BB, Melvin JL, Fiedler IG. Specificity of ICD-9-CM coding practices for stroke rehabilitation. *American Journal of Physical Medicine and Rehabilitation*. 1993;72(5):318-324.

867. Stineman MG, Hamilton BB, Goin JE, Granger CV, Fiedler RC. Functional gain and length of stay for major rehabilitation impairment categories. Patterns revealed by function related groups. *American Journal of Physical Medicine and Rehabilitation*. 1996;75(1):68-78.
868. Stineman MG, Hamilton BB, Granger CV, Goin JE, Escarce JJ, Williams SV. Four methods for characterizing disability in the formation of function related groups. *Archives of Physical Medicine and Rehabilitation*. 1994;75(12):1277-1283.
869. Stineman MG, Jette A, Fiedler RC, Granger CV. Impairment-specific dimensions within the Functional Independence Measure. *Archives of Physical Medicine and Rehabilitation*. 1997;78(6):636-643.
870. Stineman MG, Ross RN, Fielder RC, Granger CV, Maslin G. Functional independence staging: conceptual foundation, face validity, and empirical derivation. *Archives of Physical Medicine and Rehabilitation*. 2003;84(1): 29-37.
871. Stineman MG, Ross RN, Granger CV, Maislin G. Predicting the achievement of 6 grades of physical independence from data routinely collected at admission to rehabilitation. *Arch Phys Med Rehabil*. Nov 2003;84(11):1647-1656.
872. Stineman MG, Marino RJ, Deutsch A, Granger CV, Maislin G. A functional strategy for classifying patients after traumatic spinal cord injury. *Spinal Cord*. Oct 1999;37(10):717-725.
873. Stineman MG, Shea JD, Jette A, et al. The Functional Independence Measure: tests of scaling assumptions, structure, and reliability across 20 diverse impairment categories. *Archives of Physical Medicine and Rehabilitation*. 1996;77(11):1101-8.
874. Stineman MG, Tassoni CJ, Escarce JJ, et al. Development of Function-Related Groups Version 2.0: a classification system for medical rehabilitation. *Health Services Research*. 1997;32(4):529-48.
875. Stineman MG, Wechsler B, Ross R, Maislin G. A method for measuring quality of life through subjective weighting of functional status. *Arch Phys Med Rehabil*. 2003;84(4 Suppl 2):S15-22.
876. Streppel KR, Van Harten WH. The Functional Independence Measure used in a Dutch rehabilitation stroke population; a pilot study to assess progress. *Int J Rehabil Res*. 2002;25(2):87-91.
877. Suchy Y, Blint A, Osment DC. Behavioral Dyscontrol Scale: criterion and predictive validity in an inpatient rehabilitation unit population. *Clinical Neuropsychologist*. 1997;11(3):258-265.
878. Suhr JA, Grace J. Brief cognitive screening of right hemisphere stroke: relation to functional outcome. *Arch Phys Med Rehabil*. 1999;80(7):773-776.
879. Sumida M, Fujimoto M, Tokuhira A, Tominaga T, Magara A, Uchida R. Early rehabilitation effect for traumatic spinal cord injury. *Arch Phys Med Rehabil*. 2001;82(3):391-395.

880. Suzuki T, Sonoda S, Misawa K, Saitoh E, Shimizu Y, Kotake T. Incidence and consequence of falls in inpatient rehabilitation of stroke patients. *Exp Aging Res*. 2005;31(4):457-469.
881. Svensson S, Sonn U, Sunnerhagen KS. Reliability and validity of the Northwick Park Dependency Score (NPDS) Swedish version 6.0. *Clin Rehabil*. 2005;19(4):419-425.
882. Swaine BR, Pless IB, Friedman DS, Montes JL. Effectiveness of a head injury program for children: a preliminary investigation. *Am J Phys Med Rehabil*. 2000;79(5):412-420.
883. Tate DG, Forchheimer M, Daugherty J, Maynard F. Determining differences in post discharge outcomes among catastrophically and noncatastrophically sponsored outpatients with spinal cord injury. *American Journal of Physical Medicine and Rehabilitation*. 1994;73(2):89-97.
884. Tan WH, Goldstein R, Gerrard P, Ryan CM, Niewczyk P, Kowalske K, Zafonte R, Schneider JC. Outcomes and predictors in burn rehabilitation. *J Burn Care Res*. 2012;33:110-117.
885. Tanaka S, Hachisuka K, Nara S, Ogata H, Kobayashi Y, Tanaka H. Effect of activities of daily living on fiber type atrophy of the vastus lateralis muscle in patients with joint disorders. *Am J Phys Med Rehabil*. 1998;77(2):122-127.
886. Teasell R, McRae M, Foley N, Bhardwaj A. The incidence and consequences of falls in stroke patients during inpatient rehabilitation: factors associated with high risk. *Arch Phys Med Rehabil*. 2002;83(3):329-333.
887. Teasell RW, Foley NC, Bhogal SK, Chakraverty R, Bluvol A. A rehabilitation program for patients recovering from severe stroke. *Can J Neurol Sci*. 2005;32(4):512-517.
888. Teasel RW, McRae MP, Foley N, Bhardwaj A. Physical and functional correlations of ankle-foot orthosis use in the rehabilitation of stroke patients. *Arch Phys Med Rehabil*. 2001;82(8):1047-1049.
889. Teixeira da Cunha Filho I, Lim PA, Qureshy H, Henson H, Monga T, Protas EJ. A comparison of regular rehabilitation and regular rehabilitation with supported treadmill ambulation training for acute stroke patients. *J Rehabil Res Deve*. 2001;38(2):245-255.
890. Tennant A, Penta M, Tesio L, et al. Assessing and adjusting for cross-cultural validity of impairment and activity limitation scales through differential item functioning within the framework of the Rasch model: the PRO-ESOR project. *Med Care*. 2004;42(1 Suppl):137-48.
891. ter Steeg AM, Lankhorst GJ. Screening instruments for disability. *Critical Reviews in Physical and Rehabilitation Medicine*. 1994;6(1):101-112.
892. Tepper S, Beatty P, DeJong G. Outcomes in traumatic brain injury: self-report versus report of significant others. *Brain Injury*. 1996;10(8):575-581.



893. Tesio L, Franchignoni FP, Perucca L, Porta GL. The influence of age on length of stay, functional independence and discharge destination of rehabilitation inpatients in Italy. *Disabil Rehabil.* 1996;18(10):502-508.
894. Tesio L, Cantagallo A. The functional assessment measure (FAM) in closed traumatic brain injury outpatients: a Rasch-based psychometric study. *J Outcome Meas.* 1998;2(2):79-96.
895. Tesio L, Granger CV, Perucca L, Franchignoni FP, Battaglia MA, Russell CF. The FIM instrument in the United States and Italy: a comparative study. *Am J Phys Med Rehabil.* Mar 2002;81(3):168-176.
896. Tesio L, Perucca L, Battaglia MA, Franchignoni E. Quality assessment of the FIM (Functional Independence Measure) ratings through Rasch analysis. *Europa Medicophysica.* 1997;33(2):69-78.
897. Tesio L. Functional assessment in rehabilitative medicine: principles and methods. *Euro Medicophys.* 2007;43(4):515-23.
898. Thomas KS, Hicks JJ, Johnson OA. A pilot project for group cognitive retraining with elderly stroke patients. *Physical and Occupational Therapy in Geriatrics.* 1994;12(4):51-66.
899. Thornton H, Jackson D, Tuner-Stokes L. Accuracy of prediction of walking for young stroke patients by use of the FIM. *Physiother Res Int.* 2001;6(1):1-14.
900. Tofil S, Clinchot DM. Recovery of automatic and cognitive functions in traumatic brain injury using the functional independence measure. *Brain Inj.* 1996;10(12):901-910.
901. Tooth L, McKenna K, Geraghty T. Rehabilitation outcomes in traumatic spinal cord injury in Australia: functional status, length of stay and discharge setting. *Spinal Cord.* 2003;41(4):220-230.
902. Tooth LR, Ottenbacher KJ, Smith PM, et al. Effect of functional gain on satisfaction with medical rehabilitation after stroke. *Am J Phys Med Rehabil.* Sep 2003;82(9):692-699; quiz 700-691, 715.
903. Tornese D, Mele G, Respizzi S, Di Dominica F. The measure of disability among orthopedic-traumatological inpatients. *Europa Medicophysica.* 1996;32(3):145-151.
904. Toschlog EA, MacElligot J, Sagraves SG, et al. The relationship of Injury Severity Score and Glasgow Coma Score to rehabilitative potential in patients suffering traumatic brain injury. *Am Surg.* 2003;69(6):491-497; discussion 497-498.
905. Tran PV, Schwarz J, Gorman M, Helme RD. Validation of an automated up-timer for measurement of mobility in older adults. *Medical Journal of Australia.* 1997;167(8):434-436.
906. Treger I, Fromm P, Ring H, Friedman G. Association between apolipoprotein E4 and rehabilitation outcome in hospitalized ischemic stroke patients. *Arch Phys Med Rehabil.* 2003;84(7):973-976.

907. Treger I, Streifler JY, Ring H. The relationship between mean flow velocity and functional and neurologic parameters of ischemic stroke patients undergoing rehabilitation. *Arch Phys Med Rehabil.* 2005;86(3):427-430.
908. Tsang YL, Mak MK. Sit-and-reach test can predict mobility of patients recovering from acute stroke. *Arch Phys Med Rehabil.* 2004;85(1):94-98.
909. Tsuji T, Liu M, Hase K, et al. Physical fitness in persons with hemiparetic stroke: its structure and longitudinal changes during an inpatient rehabilitation programme. *Clin Rehabil.* 2004;18(4):450-460.
910. Tsuji T, Liu M, Sonoda S, Domen K, Chino N. The stroke impairment assessment set: its internal consistency and predictive validity. *Arch Phys Med Rehabil.* 2000;81(7):863-868.
911. Tsuji T, Liu M, Toikawa H, Hanayama K, Sonoda S, Chino N. ADL structure for nondisabled Japanese children based on the Functional Independence Measure for Children (WeeFIM). *Am J Phys Med Rehabil.* 1999;78(3):208-212.
912. Tsuji T, Sonoda S, Domen K, Saitoh E, Liu M, Chino N. ADL structure for stroke patients based on the Functional Independence Measure. *American Journal of Physical Medicine and Rehabilitation.* 1995;74(6):432-438.
913. Tur BS, Gursel YK, Yavuzer G, Kucudeveci A, Arasil T. Rehabilitation outcome of Turkish stroke patients: in a team approach setting. *Int J Rehabil Res.* 2003;26(4):271-277.
914. Turani N, Kemiksizoglu A, Karatas M, Ozker R. Assessment of hemiplegic gait using the Wisconsin Gait Scale. *Scand J Caring Sci.* 2004;18(1):103-108.
915. Turhan N, Saracgil N, Oztop P, Bayramoglu M. Serum albumin and comorbidity relative to rehabilitation outcome in geriatric stroke, and possible links with stroke etiology. *Int J Rehabil Res.* 2006;29(1):81-85.
916. Turkalj Z, Colja-Matic S, Vlah N, Topoljak D, Pokos L, Zadavec S. [Results of rehabilitation after ischemic cerebrovascular stroke]. *Lijec Vjesn.* 1995;117(11-12):268-271.
917. Turner S, Marosszeky B, Timms I, Boyages J. Malignant spinal cord compression: a prospective evaluation. *International Journal of Radiation Oncology, Biology, Physics.* 1993;26(1):141-146.
918. Turner-Stokes L. Standardized outcome assessment in brain injury rehabilitation for younger adults. *Disabil Rehabil.* 2002;24(7):383-389.
919. Turner-Stokes L, Paul S, Williams H. Efficiency of specialist rehabilitation in reducing dependency and costs of continuing care for adults with complex acquired brain injuries. *J Neurol Neurosurg Psychiatry.* 2006;77(5):634-639.
920. Turner-Stokes L, Tonge P, Nyein K, Hunter M, Nielson S, Robinson I. The Northwick Park Dependency Score (NPDS): a measure of nursing dependency in rehabilitation. *Clin Rehabil.* 1998;12(4):304-318.

921. Turner-Stokes L, Turner-Stokes T. The use of standardized outcome measures in rehabilitation centres in the UK. *Clin Rehabil.* 1997;11(4):306-313.
922. Udekwi P, Kromhout-Schiro S, Vaslef, Baker C, Oller D. Glasgow Coma Scale score, mortality, and functional outcome in head-injured patients. *J Trauma.* 2004;56(5):1084-1089.
923. Ueda K, Toyosato A, Nomura S. A study on the effects of short-, medium- and long-term professional oral care in elderly persons requiring long-term nursing care at a chronic or maintenance stage of illness. *Gerodontology.* 2003;20(1):50-56.
924. Unsworth CA. The concept of function. *British Journal of Occupational Therapy.* 1993;56(8):287-92.
925. Unsworth C. Clients' perceptions of discharge housing decisions after stroke rehabilitation. *Am J Occup Ther.* 1996;50(3):207-216.
926. Usuda S, Mogi S, Tomita A, Suzuki S. [Caregiver burden and subjective health level in primary caregivers of stroke patients]. *Nippon Koshu Eisei Zasshi.* 1996;43(9):854-863.
927. Valach L, Selz B, Singer S. Length of stay in the rehabilitation center, the admission functional independence measure and the functional independence measure gain. *Int J Rehabil Res.* 2004;27(2):135-143.
928. Valach L, Steck GC, Roschli C, Selz B. Rehabilitation of cognitive and motor functions in younger and older patients with vascular brain damage, rheumatologic illness and in post-surgery patients. *Int J Rehabil Res.* 2002;25(2):133-136.
929. van Baalen B, Odding E, van Woensel MP, Roebroek ME. Reliability and sensitivity to change of measurement instruments used in a traumatic brain injury population. *Clin Rehabil.* 2006;20(8):686-700.
930. van de Weg FB, Kuik DJ, Lakhorst GJ. Post-stroke depression and functional outcome: a cohort study investigating the influence of depression on functional recovery from stroke. *Clin Rehabil.* 1999;13(3):268-272.
931. van der Putten JJ, Hobart JC, Freeman JA, Thompson AJ. Measuring change in disability after inpatient rehabilitation: comparison of the responsiveness of the Barthel index and the Functional Independence Measure. *J Neurol Neurosurg Psychiatry.* 1999;66(4):480-484.
932. van der Putten JJ, Stevenson VL, Playford ED, Thompson AJ. Factors affecting functional outcome in patients with nontraumatic spinal cord lesions after inpatient rehabilitation. *Neurorehabil Neural Repair.* 2001;15(2):99-104.
933. Vangel SJ, Lichtenberg PA. Mattis Dementia Rating Scale: Clinical utility and relationship with demographic variables. *Clinical Neuropsychologist.* 1995;9(3) 209-213.

934. van Wijck FM, Pandyan AD, Johnson GR, Barnes MP. Assessing motor deficits in neurological rehabilitation: patterns of instrument usage. *Neurorehabil Neural Repair*. 2001;15(1):23-30.
935. Velozo CA. Should occupational therapy choose a single functional outcome measure? *American Journal of Occupational Therapy*. 1994;48(10):946-947.
936. Verhoef M, Barf HA, Post AW, van Asbeck FW, Gooskens RH, Prevo AJ. Functional independence among young adults with spina bifida, in relation to hydrocephalus and level of lesion. *Dev Med Child Neurol*. 2006;48(2):114-119.
937. Vincent HK, Alfano AP, Lee L, Vincent SR. Sex and age effects on outcomes of total hip replacement after inpatient rehabilitation. *Arch Phys Med Rehabil*. 2006;87(4):461-467.
938. Vincent KR, Vincent HK, Lee LW, Alfano AP. Inpatient rehabilitation outcomes in primary and revision total knee arthroplasty patients. *Clin Orthop Relat Res*. 2006;446:201-207.
939. Vincent KR, Vincent HK, Lee LW, Weng J, Alfano AP. Outcomes after inpatient rehabilitation of primary and revision total hip arthroplasty. *Arch Phys Med Rehabil*. 2006;87(8):1026-1032.
940. Vogel LC, Krajci KA, Anderson CJ. Adults with pediatric-onset spinal cord injury: part 1: prevalence of medical complications. *J Spinal Cord Med*. 2002;25(2):106-116.
941. Vogel WB, Rittman M, Bradshaw P, et al. Outcomes from stroke rehabilitation in Veterans Affairs rehabilitation units: detecting and correcting for selection bias. *J Rehabil Res Dev*. 2002;39(3):367-383.
942. Voll R, Krumm B, Schweisthal B. Functional independence measure (FIM) assessing outcome in medical rehabilitation of neurologically ill adolescents. *International Journal of Rehabilitation Research*. 2001;24(2):123-31.
943. Wade DT. *Measurement in Neurological Rehabilitation*. Oxford: Oxford University Press, 1992:218-223.
944. Wagner KA. Outcome analysis in comprehensive medical rehabilitation. In: Fuhrer MJ, ed. *Rehabilitation Outcomes: Analysis and Measurement*. Baltimore: Brookes. 1987:19-28.
945. Wagner MT, Cushman LA. Functional significance of the subcortical vascular syndrome in the rehabilitative setting. *Archives of Physical Medicine and Rehabilitation*. 1994;75(2):193-197.
946. Wagner RH, Cifu DX, Keyser-Marcus L. Functional outcome of individuals with traumatic brain injury and lower extremity deep venous thrombosis. *J Head Trauma Rehabil*. 1999;14(6):558-566.
947. Walker WC, Keyser-Marcus LA, Cifu DX, Chaudhri M. Inpatient interdisciplinary rehabilitation after total hip arthroplasty surgery: a comparison of revision and primary total hip arthroplasty. *Arch Phys Med Rehabil*. 2001;82(1):129-133.

948. Wallace D, Duncan PW, Lai SM. Comparison of the responsiveness of the Barthel Index and the motor component of the Functional Independence Measure in stroke: the impact of using different methods for measuring responsiveness. *J Clin Epidemiol*. 2002;55(9):922-928.
949. Ward I, Pivko S, Brooks G, Parkin K. Validity of the stroke rehabilitation assessment of movement scale in acute rehabilitation: a comparison with the functional independence measure and stroke impact scale-16. *PM&R*. 2011;3(11):1013-21.
950. Warschausky S, Kay JB, Kewman DG. Hierarchical linear modeling of FIM instrument growth curve characteristics after spinal cord injury. *Arch Phys Med Rehabil*. 2001;82(3):329-334.
951. Watanabe TK, O'Dell MW, Togliatti TJ. Diagnosis and rehabilitation strategies for patients with hysterical hemiparesis: a report of four cases. *Arch Phys Med Rehabil*. 1998;79(6):709-714.
952. Waters RL, Adkins RH. Firearm versus motor vehicle related spinal cord injury: preinjury factors, injury characteristics, and initial outcome comparisons among ethnically diverse groups. *Arch Phys Med Rehabil*. 1997;78(2):150-155.
953. Waters RL, Adkins RH, Sie I, Cressy J. Postrehabilitation outcomes after spinal cord injury caused by firearms and motor vehicle crash among ethnically diverse groups. *Arch Phys Med Rehabil*. 1998;79(10):1237-1243.
954. Watson AH, Kanny EM, White DM, Anson DK. Use of standardized activities of daily living rating scales in spinal cord injury and disease services. *American Journal of Occupational Therapy*. 1995;49(3):229-234.
955. Ween JE, Alexander MP, D'Esposito M, Roberts M. Incontinence after stroke in a rehabilitation setting: outcome associations and predictive factors. *Neurology*. 1996;47(3):659-663.
956. Ween JE, Alexander MP, D'Esposito M, Roberts M. Factors predictive of stroke outcome in a rehabilitation setting. *Neurology*. 1996;47(2):388-392.
957. Ween JE, Meronff ST, Alexander MP. Recovery rates after stroke and their impact on outcome prediction. *Neurorehabil Neural Repair*. 2000;14(3):229-235.
958. Weh L, Ramb JF. [Functional Independence Measure as a predictor of expected rehabilitation outcome in patients with total endoprosthesis replacement and after apoplectic infarct]. *Z Orthop Ihre Grenzgeb*. 1992;130(4):333-338.
959. Weiner DK, Bongiorno DR, Studenski SA, Duncan PW, Kochersberger GG. Does functional reach improve with rehabilitation? *Archives of Physical Medicine and Rehabilitation*. 1993;74(8):796-800.
960. Weitzenkamp DA, Whiteneck GG, Lammertse DP. Predictors of personal care assistance for people with spinal cord injury. *Arch Phys Med Rehabil*. 2002;83(10):1399-1405.

961. Wells JD, Nicosia S. Scoring acute spinal cord injury: a study of the utility and limitations of five different grading systems. *Journal of Spinal Cord Medicine*. 1995;18(1):33-41.
962. Werner RA. Predicting outcome after acute stroke with the Functional Independence Measure. *Topics in Stroke Rehabilitation*. 1994;1(3):30-39.
963. Werner RA, Kessler S. Effectiveness of an intensive outpatient rehabilitation program for postacute stroke patients. *American Journal of Physical Medicine and Rehabilitation*. 1996;75(2):114-20.
964. White MJ, Rintala DH, Hart KA, Fuhrer MJ. Sexual activities, concerns and interests of women with spinal cord injury living in the community. *American Journal of Physical Medicine and Rehabilitation*. 1993;72(6):372-378.
965. Whiteneck G, Brooks CA, Mellick D, Harrison-Felix C, Terrill MS, Noble K. Population-based estimates of outcome after hospitalization for traumatic brain injury in Colorado. *Arch Phys Med Rehabil*. 2004;85(4 Suppl 2):S73-81.
966. Whitlock JA. Functional outcome of low-level traumatically brain injured admitted to an acute care rehabilitation programme. *Brain Injury*. 1992;6(5):447-459.
967. Whitlock JA, Hamilton BB. Functional outcome after rehabilitation of severe traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 1995;76(12):1103-1112.
968. Wiart L, Come AB, Debelleix X, et al. Unilateral neglect syndrome rehabilitation by trunk rotation and scanning training. *Arch Phys Med Rehabil*. 1997;78(4):424-429.
969. Wikander B, Ekelund P, Milson I. An evaluation of multidisciplinary intervention governed by functional independence measure (FIMsm) in incontinent stroke patients. *Scand J Rehabil Med*. 1998;30(1):15-21.
970. Wilkerson DL, Batavia AI, DeJong G. Use of functional status measures for payment of medical rehabilitation services. *Archives of Physical Medicine and Rehabilitation*. 1992;73:111-120.
971. Wilkerson DL, McDermott M. Critical functional and process pathways in medical rehabilitation - outcome-to-process analysis. *Archives of Physical Medicine and Rehabilitation*. 1994;75(6):721.
972. Williams BC, Li Y, Fries BE, Warren RL. Predicting patient scores between the functional independence measure and the minimum data set: development and performance of a FIM-MDS "crosswalk". *Arch Phys Med Rehabil*. 1997;78(1):48-54.
973. Williams BK, Galea MP, Winter AT. What is the functional outcome for the upper limb after stroke? *Aust J Physiother*. 2001;47(1):19-27.
974. Williams G, Robertson V, Greenwood K. Measuring high-level mobility after traumatic brain injury. *Am J Phys Med Rehabil*. 2004;83(12):910-920.
975. Williams G, Robertson V, Greenwood K, Goldie P, Morris ME. The concurrent validity and responsiveness of the high-level mobility assessment tool for measuring the mobility

limitations of people with traumatic brain injury. *Arch Phys Med Rehabil*. 2006;87(3):437-442.

976. Wilson DB, Houle DM, Keith RA. Stroke rehabilitation: a model predicting home return. *Western Journal of Medicine*. 1991;154(5):587-590.
977. Wolfson AM, Doctor JN, Burns SP. Clinician judgments of functional outcomes: how bias and perceived accuracy affect rating. *Arch Phys Med Rehabil*. 2000;81(12):1567-1574.
978. Wright BD, Linacre JM. Observations are always ordinal; measurements, however, must be interval. *Archives of Physical Medicine and Rehabilitation*. 1989;70(12):857-860.
979. Wright BD, Linacre JM, Smith RM, Heinemann AW, Granger CV. FIM measurement properties and Rasch model details [letter]. *Scandinavian Journal of Rehabilitation Medicine*. 1997;29(4):267-272.
980. Wright BD, Linacre JM, Smith RM, Heinemann AW, Granger CV. FIM measurement properties and Rasch model details [letter; comment]. *Scandinavian Journal of Rehabilitation Medicine*. 1996;28(3):159-62.
981. Wu J, Baguley IJ. Urinary retention in a general rehabilitation unit: prevalence, clinical outcome, and the role of screening. *Arch Phys Med Rehabil*. 2005;86(9):1772-1777.
982. Yablon SA, Rock WA, Jr., Nick TG, Sherer M, McGrath CM, Goodson KH. Deep vein thrombosis: prevalence and risk factors in rehabilitation admissions with brain injury. *Neurology*. 2004;63(3):485-491.
983. Yagura H, Hatakenaka M, Miyai I. Does therapeutic facilitation add to locomotor outcome of body-weight – supported treadmill training in nonambulatory patients with stroke? A randomized controlled trial. *Arch Phys Med Rehabil*. 2006;87(4):529-535.
984. Yamada S, Liu M, Hase K, et al. Development of a short version of the motor FIM for use in long-term care settings. *J Rehabil Med*. 2006;38(1):50-56.
985. Yamakawa Y, Satoh S, Sawa S, Ohta H, Asada T. Efficacy of milnacipran on poststroke depression on inpatient rehabilitation. *Psychiatry Clin Neurosci*. 2005;59(6):705-710.
986. Yavuz N, Tezyurek M, Akyuz M. A comparison of two functional tests in quadriplegia: the quadriplegia index of function and the functional independence measure. *Spinal Cord*. 1998;36(12):832-837.
987. Yavuzer G, Kucudeveci A, Asril T, Elhan A. Rehabilitation of stroke patients: clinical profile and functional outcome. *Am J Phys Med Rehabil*. 2001;80(4):250-255.
988. Yelnik A, Bonan I, Debray M, Lo E, Gelbert F, Bussel B. Changes in the execution of a complex manual task after ipsilateral ischemic cerebral hemispheric stroke. *Arch Phys Med Rehabil*. 1996;77(8):806-810.

989. Yilmaz F, Sahin F, Aktug S, Kuran B, Yilmaz A. Long-term follow-up of patients with spinal cord injury. *Neurorehabil Neural Repair*. 2005;19(4):332-337.
990. Yokoyama O, Sakuma F, Itoh R, Sashika H. Paraplegia after aortic aneurysm repair versus traumatic spinal cord injury: functional outcome, complications, and therapy intensity off inpatient rehabilitation. *Arch Phys Med Rehabil*. Sep 2006;87(9):1189-1194.
991. Yoon SJ, Lee JH, Kim ST, Chun MH. Evaluation of traumatic brain injured patients in correlation with functional status by localized 1H-MR spectroscopy. *Clin Rehabil*. 2005;19(2):209-215.
992. Yu DT, Chae J, Walker ME, Fang ZP. Percutaneous intramuscular neuromuscular electric stimulation for the treatment of shoulder subluxation and pain in patients with chronic hemiplegia: a pilot study. *Arch Phys Med Rehabil*. 2001;82(1):20-25.
993. Zafonte RD, Hammond FM, Mann NR, Wood DL, Black KL, Millis SR. Relationship between Glasgow coma scale and functional outcome. *Am J Phys Med Rehabil*. 1996;75(5):364-369.
994. Zafonte RD, Mann NR, Millis SR, Black KL, Wood DL, Hammond F. Posttraumatic amnesia: its relation to functional outcome. *Arch Phys Med Rehabil*. 1997;78(10):1103-1106.
995. Zafonte RD, Mann NR, Millis SR, Wood DL, Lee CY, Black KL. Functional outcome after violence related traumatic injury. *Brain Inj*. 1997;11(6):403-407.
996. Zafonte RD, Ricker JH, Hanks RA, Wood DL, Amen A, Lombard L. Spinal epidural abscess: study of early outcome. *J Spinal Cord Med*. 2003;26(4):345-351.
997. Zafonte RD, Wood DL, Harrison-Felix CL, Millis SR, Valona NV. Severe penetrating head injury: a study of outcomes. *Arch Phys Med Rehabil*. 2001;82(3):306-310.
998. Zalewski C, Keller B, Bowers C, Miske P, et al. Depressive symptomatology and post-stroke rehabilitation outcome. *Clinical Gerontologist*. 1994;14(4):62-67.
999. Zamir D, Zamir M, Reitlat T, Zeev W, Polishchuk I. Readmissions to hospital within 30 days of discharge from the internal medicine wards in southern Israel. *Eur J Intern Med*. 2006;17(1):20-23.
1000. Zuhosky JP, Dugan SA, Young JL, Bode RK, Kelly JP. A retrospective review of the incidence and rehabilitation outcome of concomitant traumatic brain injury and ligamentous knee injury. *Arch Phys Med Rehabil*. 1998;79(7):805-810.
1001. Zwecker M, Levenkrohn S, Fleisig Y, Zeilig G, Ohry A, Adunsky A. Mini-Mental State Examination, cognitive FIM instrument, and the Loewenstein Occupational Therapy Cognitive Assessment: relation to function outcome of stroke patients. *Arch Phys Med Rehabil*. 2002;83(3):342-345.